HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Venlafaxine Extended-Release Tablets safely and effectively. See full prescribing information for Venlafaxine Extended-Release Tablets.

Initial U.S. Approval: 1993

Warning and Precautions (5.18)

WARNING: Suicidality and Antidepressants See full prescribing information for complete boxed warning.

Increased risk of suicidal thinking and behavior in children, adolescents and young adults taking antidepressant for major depressive disorder (MDD) and other psychiatric disorders. Venlafaxine extended-release tablets are no approved for use in pediatric patients. (5.1)

-RECENT MAJOR CHANGES-

-INDICATIONS AND USAGE--Venlafaxine extended-release tablets are a selective serotonin and norepinephrine reuptake inhibitor (SNRI) indicated for:

Maior Depressive Disorder (MDD) (1.1) Social Anxiety Disorder (SAD) (1.2)

-DOSAGE AND ADMINISTRATION--Initial Treatment (2.1)

Indication	Starting Dose	Dose Increase	Maximum Dose
Major Depressive Disorder	75 mg/day (in some patients, 37.5 mg/day for 4-7 days)	75 mg/day increments at intervals of 4 days or longer	225 mg/day
Social Anxiety Disorder	75 mg/day	No benefit at higher doses	75 mg/day

- · Venlafaxine extended-release tablets should be taken as a single daily dose with food in either the morning or evening at the
- . Discontinuation: Gradual; individualized as necessary. (2.4)
- ---DOSAGE FORMS AND STRENGTHS 37.5 mg, 75 mg, 150 mg, and 225 mg tablets (3)
- -- CONTRAINDICATIONS
- Serotonin Syndrome and MAOIs: Do not use MAOI's intended to treat psychiatric disorders with venlafaxine extended-release tablets or within 7 days of stopping treatment with venlafaxine extended-release tablets. Do not use venlafaxine extended-release release tablets within 14 days of stopping an MAOI intended to treat psychiatric disorders. In addition, do not start venlafaxine extended-release tablets in a patient who is being treated with linezolid or intravenous methylene blue (4.1).

----WARNINGS AND PRECAUTIONS----

- Serotonin Syndrome: Sertotonin syndrome has been reported with SSRIs and SNRIs, including venlafaxine extended-release tablets, both when taken alone, but especially when co-administered with other serotonergic agents (including triptans, tricyclic tables, both white hash above, our september of the commission of the second of the second of the commission of the comm extended-release tablets with other serotonergic drugs is clinically warranted, patients should be made aware of a potential ncreased risk for serotonin syndrome, particularly during treatment initiation and dose increases. (5.2).
- Suicidality: Monitor for clinical worsening and suicide risk. (5.1)
- Sustained hypertension may occur. Blood pressure monitoring recommended. (5.3)

FULL PRESCRIBING INFORMATION: CONTENTS*

- INDICATIONS AND USAGE
- 2.1 Initial Treatment

- 2.5 Switching Patients from Venlafaxine Hydrochloride Immediate-Release Tablets
- 3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 5 WARNINGS AND PRECAUTIONS

- 5.7 Changes in Weight
- 5.8 Changes in Height
- 5.11 Hyponatremia
- 5.12 Seizures
- 5.15 Interstitial Lung Disease and Eosinophilic Pneumonia
- 5.17 Laboratory Tests
- 6 ADVERSE REACTIONS
- 6.1 Clinical Studies Experience
- 7 DRUG INTERACTIONS
- 7.1 Alcohol
- 7.3 Diazepam 7.4 Haloperidol
- 7.5 Lithium 7.6 Drugs Highly Bound to Plasma Proteins

- . Angle Closure Glaucoma: Angle closure glaucoma has occurred in patients with untreated anatomically narrow angles treated
- possible). Dose reduction recommended to be gradual. (5.5)

8/2021

- Seizures have been reported. Use with caution in patients with seizure history. (5.12)
 Abnormal bleeding (most commonly ecchymosis) has been reported. (5.13)
- Interstitial lung disease and eosinophilic pneumonia have been reported. (5.15) • Sexual Dysfunction: Venlafaxine extended-release tablets may cause symptoms of sexual dysfunction. (5.18)

extended-release capsules and at a rate at least twice that of the placebo group were abnormal ejaculation, gastrointestraid complaints (nausea, dry mouth, and anorexia), CNS complaints (dizziness, somnolence, and abnormal dreams), and sweating. Social Anxiety Disorder-Adverse events in short-term studies that occurred in at least 5% of the patients receiving venlafaxine extended-release capsules and at a rate at least twice that of the placebo group were asthenia, gastrointestinal complaints (anorexia, dry mouth, nausea), CNS complaints (anxiety, insomnia, libido decreased, nervousness, somnolence, dizziness), abnormalities of sexual function (abnormal ejaculation, orgasmic dysfunction, impotence), yawn, sweating, and abnormal vision.

To report SUSPECTED ADVERSE REACTIONS, contact Camber Pharmaceuticals, Inc., at 1-866-495-8330 or FDA at 1-800-

- -- DRUG INTERACTIONS-• MAOI's: concomitant use contraindicated (4). Avoid MAOI's 14 days before starting venlafaxine and 7 days after stopping
- Cimetidine: Caution in patients with pre-existing hypertension, in elderly patients and patients with hepatic dysfunction. (7.2)
- substances that inhibit both CYP2D6 and CYP3A4. (7.7)
- Metoproloi: Possibly reduced blood-pressure lowering effect despite increased metoprolol plasma levels. Caution should be exercised with co-administration of venlafaxine and metoprolol. (7.8)
- Tryptophan supplements: Concomitant use not recommended. (7.10)
 Tryptophan supplements: Concomitant use not recommended. (7.10)
 Tryptophan supplements: Concomitant use not recommended. (7.10)
- the third trimester should be carefully considered. (2.3; 8.1)
- Nursing: Potential for serious adverse reactions in the infant. Discontinue nursing or drug, considering the importance of the
- Pediatric use: Not approved for use in pediatric patients. When considering use in a child or adolescent, balance potential risks with clinical need. (8.4)
- Hemodialysis: Reduction of daily dose by 50%. (2.3; 8.7)

Revised: 10/21

- 1.1 Major Depressive Disorde 1.2 Social Anxiety Disorder
- 2 DOSAGE AND ADMINISTRATION
- 2.2 Maintenance Treatment 2.3 Special Populations
- 2.4 Discontinuing Venlafaxine Extended-Release Tablets
- 2.6 Switching a Patient To or From a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders 2.7 Use of Venlafaxine Extended-Release Tablets with Other MAOIs, Such as Linezolid or Methylene Blue

- 5.1 Clinical Worsening and Suicide Risk 5.2 Serotonin Syndrome
- 5.3 Sustained Hypertension
- 5.4 Angle Closure Glaucoma
 5.5 Discontinuation of Treatment with Venlafaxine Extended-Release Tablets 5.6 Insomnia and Nervousness
- 5.9 Changes in Appetite
- 5.10 Activation of Mania/Hypomania
- 5.13 Abnormal Bleeding 5.14 Serum Cholesterol Elevation
- 5.16 Use in Patients With Heart Disease
- 5.18 Sexual Dysfunction
- 6.2 Post-Marketing Experience
- 7.2 Cimetidine
- 7.7 Drugs that Inhibit Cytochrome P450 Isoenzymes

- 7.8 Drugs Metabolized by Cytochrome P450 Isoenzymes 7.9 Monoamine Oxidase Inhibitors (MAOIs) 7.10 Serotonergic Drugs
- 7.11 Drugs that Interfere with Hemostasis (e.g., NSAID's, Aspirin, and Warfarin) 7.11 Didgs that interior with hemostasis (e.g., NoAid S, A. 7.12 Electroconvulsive Therapy 7.13 Postmarketing Spontaneous Drug Interaction Reports
- 7.14 Drug-Laboratory Test Interactions
- 8 USE IN SPECIFIC POPULATIONS
- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers 8.4 Pediatric Use 8.5 Geriatric Use
- 8.6 Patients with Hepatic Impairment
- 9 DRUG ABUSE AND DEPENDENCE
- 9.1 Controlled Substance 9.2 Abuse
- 9.3 Dependence 10 OVERDOSAGE
- 10.1 Human Experience
- 10.2 Management of Overdosage 11 DESCRIPTION
- 12 CLINICAL PHARMACOLOGY 12.1 Mechanism of Action
- 12 2 Pharmacodynamics
- 13 NONCLINICAL TOXICOLOGY 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES
- 14.1 Major Depressive Disorder 14.2 Social Anxiety Disorder (Social Phobia)
- 6 HOW SUPPLIED/STORAGE AND HANDLING 17 PATIENT COUNSELING INFORMATION

intervals. Individualization of tapering may be necessary.

- 17.1 Clinical Worsening and Suicide Risk 17.2 Interference with Cognitive and Motor Performance
- 17.3 Concomitant Medication 17.4 Alcohol
- 17.5 Allergic Reactions 17.6 Pregnancy
- 17.7 Nursing 17.8 Angle Closure Glaucoma 17.9 Sexual Dysfunction

3 DOSAGE FORMS AND STRENGTHS

and Warnings and Precautions (5.2).

5.1 Clinical Worsening and Suicide Risk

5 WARNINGS AND PRECAUTIONS

Age Range

<18

18-24

25-64

≥65

presenting symptoms.

4 CONTRAINDICATIONS

axine extended-release tablets are available as:

* Sections or subsections omitted from the full prescribing information are not listed.

have been reported [see Warnings and Precautions (5.5)]. Patients should be monitored for these symptoms when discontinuing

treatment. A gradual reduction in the dose rather than abrupt cessation is recommended whenever possible. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed

dose may be considered. Subsequently, the physician may continue decreasing the dose but at a more gradual rate. In clinical trials

with venlafaxine hydrochloride extended-release capsules, tapering was achieved by reducing the daily dose by 75 mg at 1 week

Depressed patients who are currently being treated at a therapeutic dose with venlafaxine hydrochloride immediate-release tablets

may be switched to venlafaxine extended-release tablets at the nearest equivalent dose (mg/day), e.g., 37.5 mg venlafaxine two-times-a-day to 75 mg venlafaxine extended-release tablets once daily. However, individual dosage adjustments may be necessary.

At least 14 days should elapse between discontinuation of an MAOI intended to treat psychiatric disorders and initiation of therapy

with venlafaxine extended-release tablets. Conversely, at least 7 days should be allowed after stopping venlafaxine extended-release tablets before starting an MAOI intended to treat psychiatric disorders [see Contraindications (4.1)].

Do not start venlafaxine extended-release tablets in a patient who is being treated with linezolid or intravenous methylene blue

because there is incleased into in Sertonian Symonthe. In a patient wind requires more up gent deathers of a psychiatric condition, other interventions, including hospitalization, should be considered [see Contraindications (4.1)]. In some cases, a patient already receiving venlafaxine extended-release tablets therapy may require urgent treatment with linezolid

or intravenous methylene blue. If acceptable alternatives to linezolid or intravenous methylene blue treatment are not available and

the potential benefits of linezolid or intravenous methylene blue treatment are judged to outweigh the risks of serotonin syndrome

in a particular patient, venlafaxine extended-release tablets should be stopped promptly, and linezolid or intravenous methylene

blue can be administered. The patient should be monitored for symptoms of serotonin syndrome for 7 days or until 24 hours after the last dose of linezolid or intravenous methylene blue, whichever comes first. Therapy with venlafaxine extended-release tablets

doses much lower than 1 mg/kg with venlafaxine extended-release tablets is unclear. The clinician should, nevertheless, be aware

The use of MAOI's intended to treat psychiatric disorders with venlafaxine extended-release tablets or within 7 days of stopping

Starting ventafaxine extended-release tablets in a patient who is being treated with MAOIs such as linezolid or intravenous methylene blue is also contraindicated because of an increased risk of serotonin syndrome [see Dosage and Administration (2.7)]

Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of their depression and/

or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression

and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a

long standing concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence

of suicidality in certain patients during the early phases of treatment. Pooled analyses of short-term placebo-controlled trials of

antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18-24) with major depressive disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24;

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, obsessive compulsive disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of 9 antidepressant drugs in over 4400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials

(median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients.

There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger patients for almost all drugs studied. There were differences in absolute risk of suicidality across the different indications, with the highest

incidence in MDD. The risk differences (drug vs placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number of cases of suicidality per 1000 patients treated) are provided in

Table 1

No suicides occurred in any of the pediatric trials. There were suicides in the adult trials, but the number was not sufficient to reach

It is unknown whether the suicidality risk extends to longer-term use, i.e., beyond several months. However, there is substantial

All patients being treated with antidepressants for any indication should be monitored appropriately and observed cl

of drug therapy, or at times of dose changes, either increases or decreases.

evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the

for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course

The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for major depressive disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a

causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality. Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients

whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to

If the decision has been made to discontinue treatment, medication should be tapered, as rapidly as is feasible, but with recognition

that abrupt discontinuation can be associated with certain symptoms [see Dosage and Administration (2.5) and Warnings and

vorsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's

Drug-Placebo Difference in Number of Cases of Suicidality per

1000 Patients Treated

Increases Compared to Placebo

14 additional cases

5 additional cases

Decreases Compared to Placebo

1 fewer case

nded to treat psychiatric disorders is als

treatment with venlafaxine extended-release tablets is contraindicated because of an increased risk of serotonin syndrome. The use of venlafaxine extended-release tablets within 14 days of stopping an MAOI intended to treat psychiatric disorders is all

may be resumed 24 hours after the last dose of linezolid or intravenous methylene blue [see Warnings and Precautions (5.2)].

The risk of administering methylene blue by non-intravenous routes (such as oral tablets or by local injection) or in intraven

of the possibility of emergent symptoms of serotonin syndrome with such use [see Warnings and Precautions (5.2)].

37.5 mg tablets (White to off white, film coated, round biconvex tablets printed with "392" in black ink)

75 mg tablets (White to off white, film coated, round biconvex tablets printed with "393" in black ink)

150 mg tablets (White to off white, film coated, round biconvex tablets printed with "394" in black ink)

contraindicated. [see Dosage and Administration (2.6) and Warnings and Precautions (5.2)].

there was a reduction with antidepressants compared to placebo in adults aged 65 and older.

ecause there is increased risk of serotonin syndrome. In a patient who requires more urgent treatment of a psychiatric condition

2.6 Switching a Patient To or From a Monoamine Oxidase Inhibitor (MAOI) Intended to Treat Psychiatric Disorders

2.7 Use of Venlafaxine Extended-Release Tablets with Other MAOIs, Such as Linezolid or Methylene Blue

FULL PRESCRIBING INFORMATION

WARNING: SUICIDALITY AND ANTIDEPRESSANT DRUGS ints increased the risk compared to placebo of suicidal thinking and behavior (suicidality) in childre adolescents, and young adults in short-term studies of Major Depressive Disorder (MDD) and other psychiatric disorders. Anyone considering the use of venlafaxine extended-release tablets or any other antidepressant in a child, ent, or young adult must balance this risk with the clinical need. Short-term studies did not show an increas in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction in risk with antidepressants compared to placebo in adults aged 65 and older. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior. Families and caregivers should be advised of the need for close observation and communication with the prescriber. Venlafaxine extended-release tablets are not approved for use in pediatric

patients. [see Warnings and Precautions (5.1) and Patient Counseling Information (17.1)] 1 INDICATIONS AND USAGE

1.1 Major Depressive Disorder

Venlafaxine extended-release tablets are indicated for the treatment of major depressive disorder (MDD). Efficacy of ventafaxine in MDD was shown in both short-term trials and a longer-term trial in MDD [see Clinical Studies (14.1)]. A major depressive episode (DSM-IV) implies a prominent and relatively persistent (nearly every day for at least 2 weeks) depresses mood or the loss of interest or pleasure in nearly all activities, representing a change from previous functioning, and includes the presence of at least five of the following nine symptoms during the same two-week period: depressed mood, markedly diminished interest or pleasure in usual activities, significant change in weight and/or appetite, insomnia or hypersomnia, psychomotor agitation or retardation, increased fatigue, feelings of guilt or worthlessness, slowed thinking or impaired concentration, a suicide

attempt or suicidal ideation.

1.2 Social Anxiety Disorder

ded-release tablets are indicated for the treatment of Social Anxiety Disorder (SAD), also known as Social Phobia, Venlafaxine extended-r as defined in DSM-IV. Social Anxiety Disorder (DSM-IV) is characterized by a marked and persistent fear of 1 or more social or performance situations in which the person is exposed to unfamiliar people or to possible scrutiny by others. Exposure to the feared situation almost invariably provokes anxiety, which may approach the intensity of a panic attack. The feared situations are avoided or endured with intense anxiety or distress. The avoidance, anxious anticipation, or distress in the feared situation(s) interferes significantly with the person's normal routine, occupational or academic functioning, or social activities or relationships, or there is a marked distress about having the phobias. Lesser degrees of performance anxiety or shyness generally do not require psychopharmacological

2 DOSAGE AND ADMINISTRATION $Ven la faxine\ extended-release\ tablets\ should\ be\ administered\ in\ a\ single\ dose\ with\ food\ either\ in\ the\ morning\ or\ in\ the\ evening\ at$ approximately the same time each day. Each tablet should be swallowed whole with fluid and not divided, crushed, chewed, or

Efficacy of venlafaxine extended-release in the treatment of SAD was established in short-term SAD trials [see Clinical Studies

placed in water. 2.1 Initial Treatment Major Depressive Disorder

 $For most patients, the recommended starting dose for venlafaxine extended-release tablets is 75 \,mg/day, administered in a single$ dose. In the clinical trials establishing the efficacy of venlafaxine hydrochloride extended-release capsules in moderately depre outpatients, the initial dose of veniafaxine was 75 mg/day. For some patients, it may be desirable to start at 37.5 mg/day for 4 to 7 days, to allow new patients to adjust to the medication before increasing to 75 mg/day. While the relationship between dose and antidepressant response for veniafaxine hydrochloride extended-release capsules has not been adequately explored, patients not responding to the initial 75 mg/day dose may benefit from dose increases to a maximum of approximately 225 mg/day. Dose increases should be in increments of up to 75 mg/day, as needed, and should be made at intervals of not less than 4 days, since steady state plasma levels of venlafaxine and its major metabolites are achieved in most patients by day 4. In the clinical trials establishing efficacy, upward titration was permitted at intervals of 2 weeks or more; the average doses were about 140 to 180

It should be noted that, while the maximum recommended dose for moderately depressed outpatients is also 225 mg/day for venlafaxine hydrochloride immediate-release tablets, more severely depressed inpatients in one study of the development program for that product responded to a mean dose of 350 mg/day (range of 150 to 375 mg/day). Whether or not higher doses of venlafaxine extended-release tablets are needed for more severely depressed patients is unknown; however, the experience with venlafaxine hydrochloride extended-release capsule doses higher than 225 mg/day is very limited. [see Warnings and Precautions (5.17)]

Social Anxiety Disorder (Social Phobia) The recommended dose is 75 mg/day, administered in a single dose. There was no evidence that higher doses confer any additional benefit. [see Warnings and Precautions (5.17)]

There is no body of evidence available from controlled trials to indicate how long patients with major depressive disorder should

be treated with venlafaxine extended-release tablets.
It is generally agreed that acute episodes of major depressive disorder require several months or longer of sustained pharmacological therapy beyond response to the acute episode. In one study, in which patients responding during 8 weeks of acute treatment with veniafaxine hydrochloride extended-release capsules were assigned randomly to placebo or to the same dose of veniafaxine hydrochloride extended-release capsules (75, 150, or 225 mg/day, qAM) during 26 weeks of maintenance treatment as they had received during the acute stabilization phase, longer-term efficacy was demonstrated. A second longer-term study has demonstrated the efficacy of venlafaxine hydrochloride immediate-release tablets in maintaining a response in patients with recurrent major depressive disorder who had responded and continued to be improved during an initial 26 weeks of treatment and were then randomly assigned to placebo or venlafaxine hydrochloride immediate-release tablets for periods of up to 52 weeks on the same dose (100 to 200 mg/day, on a b.i.d. schedule) [see Clinical Studies (14]). Based on these limited data, it is not known whether or not the dose of venlafaxine extended-release tablets needed for maintenance treatment is identical to the dose needed to achieve an initial response. Patients should be periodically reassessed to determine the need for maintenance treatment and the appropriate dose for such treatment

2.3 Special Populations Treatment of Pregnant Women During the Third Trimester

Patients with Hepatic Impairment

Neonates exposed to venlafaxine hydrochloride extended-release capsules, other SNRIs, or SSRIs, late in the third trimester have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding [see Use in Specific Populations (8.1)]. When treating pregnant women with venlafaxine extended-release tablets during the third trimester, the physician should carefully consider the potential risks and benefits of treatment.

Given the decrease in clearance and increase in elimination half-life for both venlafaxine and ODV that is observed in patients with hepatic cirrhosis and mild and moderate hepatic impairment compared with normal subjects [see Use in Specific Populations (8.6) and Clinical Pharmacology (12.3]], it is recommended that the total daily dose be reduced by 50% in patients with mild to moderate

hepatic impairment. Since there was much individual variability in clearance between patients with cirrhosis, it may be necessary to reduce the dose even more than 50%, and individualization of dosing may be desirable in some patients Given the decrease in clearance for venlafaxine and the increase in elimination half-life for both venlafaxine and ODV that is observed in patients with renal impairment (GFR = 10 to 70 mL/min) compared with normal subjects [see Use in Specific Populations (8.7) and Clinical Pharmacology (12.3], it is recommended that the total daily dose be reduced by 25% to 50%.

In patients undergoing hemodialysis, it is recommended that the total daily dose be reduced by 50%. Because there was much individual variability in clearance between patients with renal impairment, individualization of dosage may be desirable in some

Elderly Patients No dose adjustment is recommended for elderly patients solely on the basis of age. As with any drug for the treatment of major depressive disorder or Social Anxiety Disorder, however, caution should be exercised in treating the elderly. When individualizing the dosage, extra care should be taken when increasing the dose.

2.4 Discontinuing Venlafaxine Extended-Release Tablets Families and caregivers of patients being treated with antidepressants for major depressive disorder or other indications Symptoms associated with discontinuation of venlafaxine hydrochloride extended-release capsules, other SNRI's, and SSRI's both psychiatric and nonpsychiatric, should be alerted about the need to monitor patients for the emergence of agitation,

4296 Package Insert for Venlafaxine Extended-Release Tablets (Ascent-Camber) 126-10-2021.indd 1

irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to health care providers. Such monitoring should include daily observation by

with antidepressants. (5.4)
Abrupt discontinuation or dose reduction: Discontinuation symptoms may occur (generally self-limiting; serious symptoms

Activation of Mania/Hypomania has occurred. (5.10) Symptomatic hyponatremia may occur. (5.11)

. Serum cholesterol: Clinically relevant cholesterol increases may occur. Cholesterol measurements should be considered during long-term therapy. (5.14)

----ADVERSE REACTIONS--Major Depressive Disorder-Adverse events in short-term studies that occurred in at least 5% of the patients receiving venlafaxine FDA-1088 or www.fda.gov/medwatch.

Haloperidol: Increase in Haloperidol AUC and C_{max}. (7.4)
 Ketoconazole: Increase in venlafaxine and O-desmethylvenlafaxine AUC and C_{max}. Caution when using venlafaxine with

CNS-active drugs: Caution when using venlafaxine with such drugs. (7.10) Serotonergic drugs (e.g., triptans, SSRIs, other SNRIs, linezolid, lithium, tramadol, or St. John's Wort): Potential for serotonin syndrome. Careful patient observation advised. (7.10)

Pregnancy: Use during pregnancy only if clearly needed. Neonates exposed to venlafaxine in the third trimester have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding. Benefits and risk of venlafaxine use in

drug to the mother. (8.3)

Hepatic impairment: Reduction of total daily dose by 50% recommended in patients with mild to moderate impairment. In patients with cirrhosis, further reduction may be necessary and dosing individualization may be desirable. (2.3; 8.6)
Renal impairment: Reduction of daily dose by 25-50% recommended. Dosing individualization may be necessary. (2.3; 8.7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved Medication Guide.

Table 3: Incidence (%) of Sustained Elevations in SDBP in Venlafaxine Hydrochloride Immediate-Release Tablet Studies		
Venlafaxine mg/day Incidence		
<100	3%	
>100 to ≤200	5%	
>200 to ≤300	7%	
>300	13%	

Prescriptions for venlafaxine extended-release tablets should be written for the smallest quantity of tablets consistent with good

A major depressive episode may be the initial presentation of bipolar disorder. It is generally believed (though not established in

controlled trials) that treating such an episode with an antidepressant alone may increase the likelihood of precipitation of a mixed/

manic episode in patients at risk for bipolar disorder. Whether any of the symptoms described above represent such a conversion is unknown. However, prior to initiating treatment with an antidepressant, patients with depressive symptoms should be adequately

screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including

a family history of suicide, bipolar disorder, and depression. It should be noted that venlafaxine extended-release tablets are not

The development of a potentially life-threatening serotonin syndrome has been reported with SNRIs and SSRIs, including

wentafaxine extended-release tablets, alone but particularly with concomitant use of other serotonergic drugs (including triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, tryptophan, busipirone, amphetamines, and St. John's Wort), and with drugs

that impair metabolism of serotonin (in particular, MAOIs, both those intended to treat psychiatric disorders and also others, such

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, delirium, and coma), autonomic

instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, and/or gastrointestinal symptoms (e.g., nausea, vomiting,

The concomitant use of venlafaxine extended-release tablets with MAOIs intended to treat psychiatric disorders is contraindicated. Venlafaxine extended-release tablets should also not be started in a patient who is being treated with MAOIs such as linezolid

or intravenous methylene blue. All reports with methylene blue that provided information on the route of administration involved

There may be circumstances when it is necessary to initiate treatment with a MAOI such as linezolid or intravenous methylene

blue in a patient taking venlafaxine extended-release tablets. Venlafaxine extended-release tablets should be discontinued before

fentanyl, lithium, tramadol, buspirone, tryptophan, amphetamines, and St. John's Wort is clinically warranted, patients should be

Treatment with venlafaxine extended-release tablets and any concomitant serotonergic agents should be discontinued immediately

5.3 Sustained Hypertension
Venlafaxine hydrochloride extended-release capsule treatment is associated with sustained hypertension (defined as treatment

An analysis for patients in venlafaxine hydrochloride immediate-release tablet studies meeting criteria for sustained hyp

emergent supine diastolic blood pressure (SDBP) ≥90 mm Hg and ≥10 mm Hg above baseline for 3 consecutive on-therapy visits)

revealed a dose-dependent increase in the incidence of sustained hypertension for immediate-release venlafaxine hydrochloride

An insufficient number of patients received mean doses of venlafaxine hydrochloride extended-release capsules over 300 mg/day

Table 2: Number (%) of Sustained Elevations in SDBP in Venlafaxine Hydrochloride Extended-Release Capsule Premarketing

Studies by Indication

Other Clinical Trials (75-225 mg/day)

5/771 (0.6)

omitant use of venlafaxine extended-release tablets with other serotonergic drugs including triptans, tricyclic anti

aware of a potential increased risk for serotonin syndrome, particularly during treatment initiation and dose increases.

nitiating treatment with the MAOI. [see Contraindications (4.1) and Dosage and Administration (2.6 and 2.7)].

ous administration in the dose range of 1 mg/kg to 8 mg/kg. No reports involved the administration of methylene blue by

patient management, in order to reduce the risk of overdose.

approved for use in treating bipolar depression.

as linezolid and intravenous methylene blue).

diarrhea). Patients should be monitored for the emergence of serotonin syndrome.

other routes (such as oral tablets or local tissue injection) or at lower doses.

if the above events occur and supportive symptomatic treatment should be initiated

to fully evaluate the incidence of sustained increases in blood pressure at these higher doses

Major Depressive Disorder (75-375 mg/day)

19/705 (3)

5.2 Serotonin Syndrome

(see Table 3).

In premarketing major depressive disorder studies, 0.7% (5/705) of the venlafaxine hydrochloride extended-release capsuletreated patients discontinued treatment because of elevated blood pressure. Among these patients, most of the blood pressure increases were in a modest range (12 to 16 mm Hg, SDBP). In other clinical studies, 0.6% (5/771) of the venlafaxine hydrochloride extended-release capsule-treated patients discontinued treatment because of elevated blood pressure. In these patients, the blood

pressure increases were modest (1 to 24 mm Hg, SDBP). Sustained increases of SDBP could have adverse consequences. Cases of elevated blood pressure requiring immediate treating immediate have been reported in post marketing experience. Pre-existing hypertension should be controlled before treatment with venlafaxine. It is recommended that patients receiving

ventafaxine extended-release tablets have regular monitoring of blood pressure. For patients who experience a sustained increase in blood pressure while receiving ventafaxine, either dose reduction or discontinuation should be considered. Elevations in Systolic and Diastolic Blood Pressure In placebo-controlled premarketing studies, there were changes in mean blood pressure (see Table 4 for mean change in supine systolic and supine diastolic blood pressure). Across most indications, a dose-related increase in supine systolic and diastolic blood pressure was evident in venlafaxine hydrochloride extended-release capsule-treated patients.

	Venlafaxine	Venlafaxine Hydrochloride Extended-Release Capsules mg/day			Placebo	
	≤	≤75		>75		
	SSBP ¹	SDBP ²	SSBP	SDBP	SSBP	SDBP
Major Depressive Disorder 8-12 weeks	-0.28	0.37	2.93	3.56	-1.08	-0.10
Other Clinical Trials 12 weeks	-0.29	-1.26	1.18	1.34	-1.96	-1.22

¹Supine Systolic Blood Pressure

5.4 Angle Closure Glaucoma

Across all clinical trials, 1.4% of patients in the venlafaxine hydrochloride extended-release capsule-treated groups experienced a ≥15 mm Hg increase in supine diastolic blood pressure with blood pressure ≥105 mm Hg compared to 0.9% of patients in the placebo groups. Similarly, 1% of patients in the venlafaxine hydrochloride extended-release capsule-treated groups experienced a ≥20 mm Hg increase in supine systolic blood pressure with blood pressure ≥180 mm Hg compared to 0.3% of patients in the placebo groups.

Angle-Closure Glaucoma: The pupillary dilation that occurs following use of many antidepressant drugs including venlafaxine may

trigger an angle closure attack in a patient with anatomically narrow angles who does not have a patent iridectomy. 5.5 Discontinuation of Treatment with Venlafaxine Extended-Release Tablets clinical trials and retrospective surveys of trials in major depressive disorder and social anxiety disorder. Abrupt discontinuatio or dose reduction of venlafaxine at various doses has been found to be associated with the appearance of new symptoms, the frequency of which increased with increased dose level and with longer duration of treatment. Reported symptoms include agitation, anorexia, anxiety, confusion, impaired coordination and balance, diarrhea, dizziness, dry mouth, dysphoric mood, asciculation, fatigue, headaches, hypomania, insomnia, nausea, nervousness, nightmares, sensory disturbances (including shock-

like electrical sensations), somnolence, sweating, tremor, vertigo, and vomiting.

extended-release capsules in major depressive disorder studies.

Inhibitors), and SSRI's Gelective Serotonin Reuptake Inhibitors), there have been spontaneous reports of adverse reactions occurring upon discontinuation of these drugs, particularly when abrupt, including the following: dysphoric mood, irritability, agitation, dizziness, sensory disturbances (e.g. paresthesias such as electric shock sensations), anxiety, confusion, headache teltrargy, emotional lability, insomnia, hypomania, tinnitus, and seizures. While these reactions are generally self-limiting, there have been reports of serious discontinuation symptoms. Patients should be monitored for these symptoms when discontinuing treatment with venlafaxine extended-release tablets. A

During marketing of venlafaxine hydrochloride extended-release capsules, other SNRI's (Serotonin and Norepinephrine Reuptake

gradual reduction in the dose rather than abrupt cessation is recommended whenever possible. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose but at a more gradual rate [see Dosage and Administration (2.4).

5.6 Insomnia and Nervousness ment-emergent insomnia and nervousness were more commonly reported for patients treated with venlafaxine hydrochloride extended-release capsules than with placebo in pooled analyses of short-term major depressive disorder and other clinical studies,

as shown in Table 5 Table 5 Incider	nce of Insomnia and Nervousness in	ed Major Depressive Disorder ar Other Trials		
	Venlafaxine Hydrochloride Extended-Release Capsules	Placebo	Venlafaxine Hydrochloride Extended-Release Capsules	Placebo
Symptom	n = 357	n = 285	N = 819	n = 695
Insomnia	17%	11%	24%	8%
Nervousness	10%	5%	10%	5%

Insomnia and nervousness each led to drug discontinuation in 0.9% of the patients treated with venlafaxine hydrochloride

In other clinical trials, insomnia and nervousness led to drug discontinuation in 2% and 1%, respectively, of the patients treated with venlafaxine hydrochloride extended-release capsules up to 12 weeks. Adult Patients: A loss of 5% or more of body weight occurred in 7% of patients treated with venlafaxine hydrochloride extendedrelease capsules and 2% of placebo-treated patients in the short-term placebo-controlled major depressive disorder trials. The discontinuation rate for weight loss associated with venlafaxine hydrochloride extended-release capsules was 0.1% in major depressive disorder studies. In other placebo-controlled trials, 4% of the patients treated with venlafaxine hydrochloride extendedrelease capsules and 1% of the placebo-treated patients sustained a loss of 7% or more of body weight during up to 6 months ent. None of the patients receiving venlafaxine hydrochloride extended-release capsules in other stud

for weight loss. The safety and efficacy of venlafaxine therapy in combination with weight loss agents, including phentermine, have not been established. Co-administration of venlafaxine extended-release tablets and weight loss agents is not recommended. Venlafaxine extended-release tablets are not indicated for weight loss alone or in combination with other products.

Pediatric Patients: Weight loss has been observed in pediatric patients (ages 6-17) receiving venlafaxine hydrochloride extended release capsules. In a pooled analysis of four eight-week, double-blind, placebo-controlled, flexible dose outpatient trials for major depressive disorder (MDD) and another disorder, patients treated with venlafaxine hydrochloride extended-release capsules lost an verage of 0.45 kg (n = 333), while placebo-treated patients gained an average of 0.77 kg (n = 333). More patie renlafaxine hydrochloride extended-release capsules than with placebo experienced a weight loss of at least 3.5% in the studies (18% of patients treated with venlafaxine hydrochloride extended-release capsules vs. 3.6% of placebo-treated patients; p<0.001). n a 16-week, double-blind, placebo-controlled, flexible dose outpatient study for another disorder, venlafaxine hydrochlor extended-release capsule-treated patients lost an average of 0.75 kg (n=137), while placebo-treated patients gained an average of 0.76 kg (n=148). More patients treated with venlafaxine hydrochloride extended-release capsules than with placebo experienced a weight loss of at least 3.5% in the study (47% of patients treated with venlafaxine hydrochloride extended-release capsules vs. 14% of placebo-treated patients; p<0.001). Weight loss was not limited to patients with treatment-emergent anorexia [see Warnings and Precautions (5.9)].

The risks associated with longer-term use of venlafaxine hydrochloride extended-release capsules were assessed in an open-label MDD study of children and adolescents who received venlafaxine hydrochloride extended-release capsules for up to six months. The children and adolescents in the study had increases in weight that were less than expected based on data from age-and sex matched peers. The difference between observed weight gain and expected weight gain was larger for children (<12 years old)

Pediatric Patients: During an eight-week, placebo-controlled non-MDD study, venlafaxine hydrochloride extended-release capsule

treated patients (ages 6-17) grew an average of 0.3 cm (n=122), while placebo-treated patients grew an average of 1.0 cm (n=132); p=0.041. This difference in height increase was most notable in patients younger than twelve. During the eight-week

placebo-controlled MDD studies, venlafaxine hydrochloride extended-release capsule-treated patients grew an average of 0.8 cm

(n = 146), while placebo-treated patients grew an average of 0.7 cm (n = 147). During a 16-week, placebo-controlled non-MDD study, both the venlafaxine hydrochloride extended-release capsule-treated patients (n=109) and the placebo-treated (n=112) patients each grew an average of 1.0 cm. In the six-month, open-label MDD study, children and adolescents had height increases that were less than expected based on data from age-and sex-matched peers. The difference between observed growth rates and expected growth rates was larger for children (<12 years old) than for adolescents (≥12 years old).

Adult Patients: Treatment-emergent anorexia was more commonly reported for patients treated with venlafaxine hydrochloride extended-release capsules (8%) than for placebo-treated patients (4%) in the pool of short-term, double-blind, placebo-controlled major depressive disorder studies. The discontinuation rate for anorexia associated with venlafaxine hydrochloride extended-release capsules was 1.0% in major depressive disorder studies. Treatment-emergent anorexia was more commonly reported for patients treated with venlafaxine hydrochloride extended-release capsules (20%) than for placebo-treated patients (2%) in the pool of short-term, double-blind, placebo-controlled Social Anxiety Disorder studies. The discontinuation rate for anorexia was 0.4% for patients receiving venlafaxine hydrochloride extended-release capsules for up to 12 weeks in Social Anxiety Disorder studies Pediatric Patients: Decreased appetite has been observed in pediatric patients receiving venlafaxine hydrochloride extended release capsules. In placebo-controlled trials in MDD and another disorder, 10% of patients aged 6-17 treated with venlafaxine hydrochloride extended-release capsules for up to eight weeks and 3% of patients treated with placebo reported treatmentemergent anorexia (decreased appetite). None of the patients receiving venlafaxine hydrochloride extended-release capsules discontinued for anorexia or weight loss.

In a placebo-controlled non-MDD trial, 22% and 3% of patients aged 8-17 treated for up to 16 weeks with venlafaxir hydrochloride extended-release capsules and placebo, respectively, reported treatment-emergent anorexia (decreased appetite). The discontinuation rates for anorexia were 0.7% and 0.0% for patients receiving venlafaxine hydrochloride extended-release capsules and placebo, respectively; the discontinuation rates for weight loss were 0.7% for patients receiving either venlafaxine hydrochloride extended-release capsules or placebo. 5.10 Activation of Mania/Hynomania

During premarketing major depressive disorder studies, mania or hypomania occurred in 0.3% of patients treated with venlafaxine hydrochloride extended-release capsules and 0.0% placeho natients. In premarketing Social Anxiety Disorder studies, no natients related with venlafaxine hydrochloride extended-release capsules and no placebo-treated patients experienced mania or hypomania. In all premarketing major depressive disorder trials with venlafaxine hydrochloride immediate-release tablets, mania or hypomania occurred in 0.5% of venlafaxine-treated patients compared with 0% of placebo patients. Mania/hypomania has also been reported in a small proportion of patients with mood disorders who were treated with other marketed drugs to treat major depressive disorder. As with all drugs effective in the treatment of major depressive disorder, venlafaxine extended-release tablets should be used cautiously in patients with a history of mania.

5.11 Hyponatremia Hyponatremia may occur as a result of treatment with SSRI's and SNRI's, including venlafaxine extended-release tablets. In many cases, this hyponatremia appears to be the result of the syndrome of inappropriate antidiuretic hormone secretion (SIADH). Cases with serum sodium lower than 110 mmol/L have been reported. Elderly patients may be at greater risk of developing hyponatremia with SSRIs and SNRIs. Also, patients taking diuretics or who are otherwise volume depleted may be at greater risk five Use in Specific Populations (8.5)]. Discontinuation of venlafaxine extended-release tablets should be considered in patients with symptomatic hypopatremia and appropriate medical intervention should be instituted

Signs and symptoms of hypomatremia include headache, difficulty concentrating, memory impairment, confusion, weakness, and unsteadiness, which may lead to falls. Signs and symptoms associated with more severe and/or acute cases have included hallucination, syncope, seizure, coma, respiratory arrest, and death. 5.12 Seizures

During premarketing experience, no seizures occurred among 705 patients treated with venlafaxine hydrochloride extended-release capsules in the major depressive disorder studies or among 277 patients treated with venlafaxine hydrochloride extended-release capsules in Social Anxiety Disorder studies. In all premarketing major depressive disorder trials with venlafaxine hydrochloride immediate-release tablets, seizures were reported at various doses in 0.3% (8/3082) of venlafaxine-treated patients. Venlafaxine extended-release tablets, like many antidepressants, should be used cautiously in patients with a history of seizures and should be discontinued in any patient who develops seizures.

5.13 Abnormal Bleeding ding venlafaxine extended-release tablets, may increase the risk of bleeding events. Concomitant use

5.14 Serum Cholesterol Elevation

5.18 Sexual Dysfunction

6 ADVERSE REACTIONS

Data Sources

6.1 Clinical Studies Experience

of aspirin, nonsteroidal anti-inflammatory drugs, warfarin, and other anticoagulants may add to this risk. Case reports and epidemiological studies (case-control and cohort design) have demonstrated an association between use of drugs that interfere with serotonin reuptake and the occurrence of gastrointestinal bleeding. Bleeding events related to SSRIs and SNRIs use have ranged from ecchymoses, hematomas, epistaxis, and petechiae to life-threatening hemorrhages. Patients should be cautioned about the risk of bleeding associated with the concomitant use of venlafaxine extended-release tablets and NSAIDs, aspirin, or other drugs that affect coagulation

treated patients treated for at least 3 months in placebo-controlled trials [see Adverse Reactions (6.1)]. Measurement of serum cholesterol levels should be considered during long-term treatment. 5.15 Interstitial Lung Disease and Eosinophilic Pneumonia Interstitial lung disease and eosinophilic pneumonia associated with venlafaxine therapy have been rarely reported. The possibility

Clinically relevant increases in serum cholesterol were recorded in 5.3% of venlafaxine-treated patients and 0.0% of placebo-

of these adverse reactions should be considered in venlafaxine-treated patients who present with progressive dyspnea, cough or omfort. Such patients should undergo a prompt medical evaluation, and discontinuation of venlafaxine therapy should

5.16 Use in Patients With Heart Disease Premarketing experience with venlafaxine in patients with concomitant systemic illness is limited. Caution is advised in administering venlafaxine extended-release tablets to patients with diseases or conditions that could affect he

Venlafaxine has not been evaluated or used to any appreciable extent in patients with a recent history of myocardial infarction or unstable heart disease. Patients with these diagnoses were systematically excluded from many clinical studies during venlafaxine's premarketing testing. The electrocardiograms were analyzed for 275 patients who received venlafaxine hydrochloride extendedrelease capsules and 220 patients who received placebo in 8 to 12 week double-blind, placebo-controlled trials in major depressive disorder as well as for 195 patients who received venlafaxine hydrochloride extended-release capsules and 228 patients who received venlafaxine hydrochloride extended-release acquises and 228 patients who received placebo in 12-week double-blind, placebo-controlled trials in Social Anxiety Disorder. The mean change from baseline in corrected QT interval (QTc) for patients treated with venlafaxine hydrochloride extended-release capsules in major depressive disorder studies was increased relative to that for placebo-treated patients (increase of 4.7 mase for venlafaxine hydrochloride extended-release capsules and decrease of 1.9 msec for venlafaxine in QTC for patients treated with venlafaxine hydrochloride extended-release capsules in the Social Anxiety Disorder studies was increased relative to that for placebo-treated patients (increase of 2.8 msec for venlafaxine hydrochloride extended-release capsules and decrease of 2.0 msec for placebo).

In these same trials, the mean change from baseline in heart rate for patients treated with venlafaxine hydrochloride extended release capsules in the major depressive disorder studies was significantly higher than that for placebo (a mean increase of 4 beats per minute for veniafaxine hydrochloride extended-release capsules and 1 beat per minute for placebo). The mean change from baseline in heart rate for patients treated with venlafaxine hydrochloride extended-release capsules in the Social Anxiety Disorder studies was significantly higher than that for placebo (a mean increase of 5 beats per minute for venlafaxine hydrochloride

In a flexible-dose study, with doses of venlafaxine hydrochloride immediate-release tablets in the range of 200 to 375 mg/day and mean dose greater than 300 mg/day, patients treated with venlafaxine hydrochloride immediate-release tablets had a mean increase in heart rate of 8.5 beats per minute compared with 1.7 beats per minute in the placebo group. As increases in heart rate were observed, caution should be exercised in patients whose underlying medical conditions might

be compromised by increases in heart rate (e.g., patients with hyperthyroidism, heart failure, or recent myocardial infarction) Evaluation of the electrocardiograms for 769 patients who received venlafaxine hydrochloride immediate-release tablets in 4 to 6 week double-blind, placebo-controlled trials showed that the incidence of trial-emergent conduction abnormalities did not differ from that with placebo. 5.17 Laboratory Tests

(6.1)]. In male patients, SNRI use may result in ejaculatory delay or failure, decreased libido, and erectile dysfunction. In female patients, SNRI use may result in decreased libido and delayed or absent orgasm. It is important for prescribers to inquire about sexual function prior to initiation of venlafaxine extended-release tablets and to inquire specifically about changes in sexual function during treatment, because sexual function may not be spontaneously reported. When evaluating changes in sexual function, obtaining a detailed history (including timing of symptom onset) is important because sexual symptoms may have other causes, including the underlying psychiatric disorder. Discuss potential management strategies to support patients in making informed decisions about treatment.

Use of SNRIs, including venlafaxine extended-release tablets, may cause symptoms of sexual dysfunction [see Adverse Reactions

extended-release capsules in the entire development program for the formulation and with ventakarine hydrochloride immediate-release tablets is included in the subsection "Other Adverse Reactions Observed During the Premarketing Evaluation of Venlafaxine Hydrochloride Immediate-Release Tablets and Venlafaxine Hydrochloride Extended-Release Capsules" [see also Warnings and Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a

The information included in subsection "Adverse Findings Observed in Short-Term, Placebo-Controlled Studies with Venlafaxine

Hydrochloride Extended-Release Capsules" is based on data from a pool of three 8 and 12 week controlled clinical trials in major depressive disorder (includes two U.S. trials and one European trial), and on data up to 12 weeks from a pool of two controlled

clinical trials in Social Anxiety Disorder, Information on additional adverse reactions associated with venlafaxine hydrochloride

Adverse Findings Observed in Short-Term, Placebo-Controlled Studies with Venlafaxine Hydrochloride Extended-Release Capsules Adverse Reactions Associated with Discontinuation of Treatment Major Depressive Disorder: Approximately 11% of the 357 patients who received venlafaxine hydrochloride extended-release capsules in placebo-controlled clinical trials for major depressive disorder discontinued treatment due to an adverse reaction, compared with 6% of the 285 placeho-treated patients in those studies. Adverse reactions that led to treatment discontinuation in

Social Anxiety Disorder: Approximately 17% of the 277 patients who received venlafaxine hydrochloride extended-release capsules

least 2% of drug-treated patients were nausea, dizziness, and somnolence.

drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

in placebo-chronical trials for Social Anxiety Disorder discontinued treatment due to an adverse reaction, compared with 5% of the 274 placebo-treated patients in those studies. Adverse reactions that led to treatment discontinuation in a least 2% of drug-treated patients were nausea, insomnia a headache dizziness and somnolence Adverse Reactions Occurring at an Incidence of 5% or More Major Depressive Disorder: Note in particular the following adverse reactions that occurred in at least 5% of the patients receiving venlafaxine hydrochloride extended-release capsules and at a rate at least twice that of the placebo group for all placebo-controlled trials for the major depressive disorder indication (see Table 6): Abnormal ejaculation, gastrointestinal complaints (nausea, dry

In the two U.S. placebo-controlled trials, the following additional reactions occurred in at least 5% of patients treated with venlafaxine hydrochloride extended-release capsules (n = 192) and at a rate at least twice that of the placebo group: Abnormalities of sexual function (impotence in men, anorgasmia in women, and libido decreased), gastrointestinal complaints (constipation and flatulence), CNS complaints (insomnia, nervousness, and tremor), problems of special senses (abnormal vision), cardiovascula effects (hypertension and vasodilatation), and yawning. Social Anxiety Disorder: Note in particular the following adverse reactions that occurred in at least 5% of the patients receiving venlataxing hydrochloride extended-release capsules and at a rate at least twice that of the placeho group for the 2 placeho ventatation hypotronidae exteriorition of section and a lateral reast whice that of the placetory group for the 2 placetory controlled trials for the Social Anxiety Disorder indication (see Table 7). Asthenia, gastrointestinal complaints (anorexia, constipation, dry mouth, nausea), CNS complaints (dizziness, insomnia, libido decreased, nervousness, somnolence), abnormalities of sexual

function (abnormal ejaculation, impotence, libido decreased, orgasmic dysfunction), yawn, sweating, and abnormal vision.

Adverse Reactions Occurring at an Incidence of 2% or More Among Patients Treated with Venlafaxine Hydrochloride Extended

Tables 6 and 7 enumerate the incidence, rounded to the nearest percent, of adverse reactions that occurred during acute therapy of major depressive disorder (up to 12 weeks; dose range of 75 to 225 mg/day) and of Social Anxiety Disorder (up to 12 weeks;

mouth, and anorexia), CNS complaints (dizziness, somnolence, and abnormal dreams), and sweating.

dose range of 75 to 225 mg/day), respectively, in 2% or more of patients treated with venlafaxine hydrochloride extended-release capsules where the incidence in patients treated with venlataxine hydrochloride extended-release capsules was greater than the incidence for the respective placebo-treated patients. The table shows the percentage of patients in each group who had at least one episode of a reaction at some time during their treatment. Reported adverse reactions were classified using a standard COSTART-based Dictionary terminology. The prescriber should be aware that these figures cannot be used to predict the incidence of adverse reactions in the cours of usual medical practice where patient characteristics and other factors differ from those which prevailed in the clinical trials

treatments, uses and investigators. The cited figures, however, do provide the prescribing physician with some basis for estimating the relative contribution of drug and nondrug factors to the adverse reaction incidence rate in the population studied. ence in Short-Term Placeho-Controlled Clinical Trials with

Similarly, the cited frequencies cannot be compared with figures obtained from other clinical investigations involving different

	% Reporting Reaction			
Body System Preferred Term	Venlafaxine Hydrochloride Extended-Release Capsules (n=357)	Placebo (n=285)		
Body as a whole	Capsules (II=337)	(11=203)		
•	8%	7%		
Asthenia	8%	7%		
Cardiovascular System				
Vasodilation ³	4%	2%		
Hypertension	4%	1%		
Digestive System				
Nausea	31%	12%		
Constipation	8%	5%		
Anorexia	8%	4%		
Vomiting	4%	2%		
Flatulence	4%	3%		
Metabolic/Nutritional				
Weight Loss	3%	0%		
Nervous System				
Dizziness	20%	9%		
Somnolence	17%	8%		
Insomnia	17%	11%		
Dry Mouth	12%	6%		
Nervousness	10%	5%		
Abnormal Dreams ⁴	7%	2%		
Tremor	5%	2%		
Depression	3%	<1%		
Paresthesia	3%	1%		
Libido Decreased	3%	<1%		
Agitation	3%	1%		
Respiratory System				
Pharyngitis	7%	6%		
Yawn	3%	0%		
Skin				
Sweating	14%	3%		
Special Senses				
Abnormal Vision ⁵	4%	<1%		
Urogenital System				
Abnormal Ejaculation (male) ^{6,7}	16%	<1%		
Impotence ⁷	4%	<1%		
Anorgasmia (female)8,9	3%	<1%		

Incidence, rounded to the nearest %, for reactions reported by at least 2% of patients treated with venlafaxine hydrochloride extended-release capsules, except for reactions which had an incidence equal to or less than placebo 2 < 1% indicates an incidence greater than zero but less than 1%.

10/19/21 12:38 PM

4 Mostly "vivid dreams," "nightmares," and "increased dreaming." 5 Mostly "blurred vision" and "difficulty focusing eyes."

⁶ Mostly "delayed ejaculation." ⁷ Incidence is based on the number of male patients.

8 Mostly "delayed orgasm" or "anorgasmia

Table 7 Treatment-Emergent Adverse Reaction Incidence in Short-Term Placebo-Controlled Clinical Trials with Venlafaxine Hydrochloride Extended-Release Capsules in Social Anxiety Disorder Patients^{1,2}

	% Reporting Reaction			
Body System Preferred Term	Venlafaxine Hydrochloride Extended-Release Capsules (n=277)	Placebo (n=274)		
Body as a whole	Capsules (II—277)	(11=274)		
Headache	34%	33%		
Asthenia	17%	8%		
Flu Syndrome	6%	5%		
Accidental Injury	5%	3%		
Abdominal Pain	4%	3%		
Cardiovascular System	470	370		
	5%	4%		
Hypertension Vasodilation ³	3%			
		1%		
Palpitation	3%	1%		
Digestive System	00%	00/		
Nausea	29%	9%		
Anorexia ⁴	20%	1%		
Constipation	8%	4%		
Diarrhea	6%	5%		
Vomiting	3%	2%		
Eructation	2%	0%		
Metabolic/Nutritional				
Weight Loss	4%	0%		
Nervous System				
Insomnia	23%	7%		
Dry Mouth	17%	4%		
Dizziness	16%	8%		
Somnolence	16%	8%		
Nervousness	11%	3%		
Libido Decreased	9%	<1%		
Anxiety	5%	3%		
Agitation	4%	1%		
Tremor	4%	<1%		
Abnormal Dreams ⁵	4%	<1%		
Paresthesia	3%	<1%		
Twitching	2%	0%		
Respiratory System				
Yawn	5%	<1%		
Sinusitis	2%	1%		
Skin				
Sweating	13%	2%		
Special Senses	.570	270		
Abnormal Vision ⁶	6%	3%		
Urogenital System	070	370		
Abnormal Ejaculation ^{7,8}	16%	1%		
Impotence ⁸	10%	1%		
•				
Orgasmic Dysfunction ^{9,10}	8%	0%		

the placebo rate are not included

2 <1% means greater than zero but less than 1%.</p>
3 Mostly "hot flashes."

4 Mostly "decreased appetite" and "loss of appetite.

⁵ Mostly "vivid dreams," "nightmares," and "increased dreaming."
⁶ Mostly "blurred vision."

Includes "delayed ejaculation" and "anorgasmia."

8 Percentage based on the number of males (venlafaxine hydrochloride extended-release capsules = 158, placebo = 153). Includes "abnormal orgasm" and "anorgasmia.' ¹⁰ Percentage based on the number of females (venlafaxine hydrochloride extended-release capsules = 119, placebo = 121).

Vital Sign Changes Treatment with venlafaxine hydrochloride extended-release capsules treatment for up to 12 weeks in premarketing placebo-

controlled major depressive disorder trials was associated with a mean final on-therapy increase in pulse rate of approximately 2 beats per minute, compared with 1 beat per minute for placebo.

Treatment with venlafaxine hydrochloride extended-release capsules for up to 12 weeks in premarketing placebo-controlled Social Anxiety Disorder trials was associated with a mean final on-therapy increase in pulse rate of approximately 4 beats per minute, compared with an increase of 1 beat per minute for placebo. [see Warnings and Precautions (5.3) for effects on blood pressure.] In a flexible-dose study in MDD, with doses of venlafaxine hydrochloride immediate-release tablets in the range of 200 to 375 mg/day and mean dose greater than 300 mg/day, the mean pulse was increased by about 2 beats per minute compared with a decrease of about 1 beat per minute for placebo. [see Warnings and Precautions (5.16) for effects on heart rate.]

Laboratory Changes Serum Cholesterol

Venlafaxine hydrochloride extended-release capsules treatment for up to 12 weeks in premarketing placebo-controlled trials for major depressive disorder was associated with a mean final on-therapy increase in serum cholesterol concentration of approximately 1.5 mg/dL compared with a mean final decrease of 7.4 mg/dL for placebo. Venlafaxine hydrochloride extendedrelease capsules treatment for up to 12 weeks in other premarketing placebo-controlled trials was associated with mean final on-therapy increases in serum cholesterol concentration of approximately 7.9 mg/dL compared with a mean final decrease of 2.9 mg/dL for placebo.

ts treated with extension trials had a mean final on-therapy increase in total cholesterol of 9.1 mg/dL compared with a decrease of 7.1 mg/dL among placebo-treated patients. This increase was duration dependent over the study period and tended to be greater with higher doses. Clinically relevant increases in serum cholesterol, defined as 1) a final on-therapy increase in serum cholesterol ≥50 mg/ dL from baseline and to a value ≥261 mg/dL, or 2) an average on-therapy increase in serum cholesterol ≥50 mg/dL from baseline and to a value ≥261 mg/dL, were recorded in 5.3% of venlafaxine-treated patients and 0.0% of placebo-treated patients [see Warnings and Precautions (5.14)1.

Serum Triglycerides chloride extended-release capsules treatment for up to 12 weeks in pooled premarketing trials was associated with a mean final on-therapy increase in fasting serum triglyceride concentration of approximately 8.2 mg/dl, compared with a mean final increase of 0.4 mg/dl for placebo.

ECG Changes In a flexible-dose MDD study with doses of venlafaxine hydrochloride immediate-release tablets in the range of 200 to 375 mg/day

and mean dose greater than 300 mg/day, the mean change in heart rate was 8.5 beats per minute compared with 1.7 beats per minute for placebo. [See Warnings and Precautions (5.16)] Other Adverse Reactions Observed During the Premarketing Evaluation of Venlafaxine Hydrochloride Immediate-Release Tablets

and Venlafaxine Hydrochloride Extended-Release Capsules During its premarketing assessment, multiple doses of venlafaxine hydrochloride extended-release capsules were administered

burning its premiarching assessment, multiple duses of verniandament rypurculonide extended tease capsules were administered to 705 patients in Phase 3 major depressive disorder studies and venlafaxine hydrochloride immediate-release tablets was administered to 96 patients. During its premarketing assessment, multiple doses of venlafaxine hydrochloride extended-release capsules were also administered to 3514 patients in other Phase 3 studies. In addition, in premarketing assessment of venlafaxine hydrochloride immediate-release tablets, multiple doses were administered to 2897 patients in Phase 2 to Phase 3 studies for major depressive disorder. The conditions and duration of exposure to venlafaxine in both development programs varied greatly and included (in overlapping categories) open and double-blind studies, uncontrolled and controlled studies, inpatient (venlafaxine hydrochloride immediate-release tablets only) and outpatient studies, fixed-dose, and titration studies. Adverse reactions associated with this exposure were recorded by clinical investigators using terminology of their own choosing. Consequently, it is not possible to provide a meaningful estimate of the proportion of individuals experiencing adverse reactions without first grouping similar types of untoward events into a smaller number of standardized reaction categories.

In the tabulations that follow, reported adverse reactions were classified using a standard COSTART-based Dictionary terminology. The frequencies presented, therefore, represent the proportion of the 7212 patients exposed to multiple doses of either formulation of venlafaxine who experienced a reaction of the type cited on at least one occasion while receiving venlafaxine. All reported reactions are included except those already listed in Tables 6 and 7 and those reactions for which a drug cause was remote. If the COSTART term for a reaction was so general as to be uninformative, it was replaced with a more informative term. It is important to emphasize that, although the reactions reported occurred during treatment with venlafaxine, they were not necessarily caused by it.

Reactions are further categorized by body system and listed in order of decreasing frequency using the following definitions: frequent adverse reactions are defined as those occurring on one or more occasions in at least 1/100 patients; infrequent adverse reactions are those occurring in 1/100 to 1/1000 patients; rare reactions are those occurring in fewer than 1/1000 patients. Body as a whole - Frequent: chest pain substernal, chills, fever, neck pain; Infrequent: face edema, intentional injury, malaise

moniliasis, neck rigidity, pelvic pain, photosensitivity reaction, suicide attempt, withdrawal syndrome; Rare: appendicitis, bacteremia, cellulitis, granuloma. Cardiovascular system - Frequent: migraine, tachycardia; Infrequent: angina pectoris, bradycardia, extrasystoles, hypoteris, bradycardia, hypoteris, hypoteri peripheral vascular disorder (mainly cold feet and/or cold hands), postural hypotension, syncope; Rare: aortic aneurysm, arteritis,

first-degree atrioventricular block, bigeminy, bundle branch block, capillary fragility, cerebral ischemia, coronary artery disease, congestive heart failure, heart arrest, hematoma, cardiovascular disorder (mitral valve and circulatory disturbance), mucocutaneous hemorrhage, myocardial infarct, pallor, sinus arrhythmia, thrombophlebitis. Digestive system - Frequent: increased appetite; Infrequent: bruxism, colitis, dysphagia, tongue edema, eructation, esophagitis

gastritis, gastrointestinal ulcer, gingivitis, glossitis, rectal hemorrhage, hemorrhoids, melena, oral moniliasis, stomatitis, mouth ulceration; Rare: abdominal distension, biliary pain, chellitis, cholecystitis, cholelithiasis, esophageal spasms, duodenitis, hematemesis, gastroesophageal reflux disease, gastrointestinal hemorrhage, gum hemorrhage, hepatitis, ileitis, jaundice, intestinal obstruction, liver tenderness, parotitis, periodontitis, proctitis, salivary gland enlargement, increased salivation

Endocrine system - Rare: galactorrhoea, goiter, hyperthyroidism, hypothyroidism, thyroid nodule, thyroiditis

Hemic and lymphatic system - Frequent: ecchymosis; Infrequent: anemia, leukocytosis, leukopenia, lymphadenopathy, thrombocythemia; Rare: basophilia, bleeding time increased, cyanosis, eosinophilia, lymphocytosis, multiple myeloma, purpura, thrombocytopenia.

Metabolic and nutritional - Frequent: edema, weight gain; Infrequent: alkaline phosphatase increased, dehydration plesteremia, hyperglycemia, hyperlipidemia, hypokalemia, SGOT (AST) increased, SGPT (ALT) increased, thirst; Rare: alcohol intolerance, bilirubinemia, BUN increased, creatinine increased, diabetes mellitus, glycosuria, gout, healing abnormal hemochromatosis, hypercalcinuria, hyperkalemia, hyperphosphatemia, hyperuricemia, hypocholest hyponatremia, hypophosphatemia, hypoproteinemia, uremia.

Musculoskeletal system - Infrequent: arthritis, arthrosis, bone spurs, bursitis, leg cramps, myasthenia, tenosynovitis; Rare: pathological fracture, muscle cramp, muscle spasms, musculoskeletal stiffness, myopathy, osteoporosis, osteosclerosis, plantar fasciitis, rheumatoid arthritis, tendon rupture.

Nervous system - Frequent: amnesia. confusion, depersonalization, hypesthesia, trismus, vertigo; Infrequent: akathisia, apathy ataxia, circumoral paresthesia, CNS stimulation, emotional lability, euphoria, hallucinations, hostility, hyperesthesia, hyperkinesia hypotonia, incoordination, libido increased, manic reaction, myoclonus, neuralgia, neuropathy, psychosis, seizure, stupor, suicidal ideation; Rare: akinesia, alcohol abuse, aphasia, bradykinesia, buccoglossal syndrome, cerebrovascular accident, feeling drunk, loss of consciousness, delusions, dementia, dystonia, energy increased, facial paralysis, abnormal gait, Guillain-Barre Syndrome, homicidal ideation, hyperchlorhydria, hypokinesia, hysteria, impulse control difficulties, motion sickness, neuritis, nystagmus, paranoid reaction, paresis, psychotic depression, reflexes decreased, reflexes increased, torticollis.

Respiratory system - Frequent: cough increased, dyspnea; Infrequent: asthma, chest congestion, epistaxis, hyperventilation laryngismus, laryngitis, pneumonia; Rare: atelectasis, hemoptysis, hypoventilation, hypoxia, larynx edema, pleurisy, pulmonary

embolus, sleep apnea. Skin and appendages - Frequent: pruritus; Infrequent: acne, alopecia, contact dermatitis, dry skin, eczema, maculopapular rash psoriasis, urticaria; Rare: brittle nails, erythema nodosum, exfoliative dermatitis, lichenoid dermatitis, furunculosis, hirsutism leukoderma, miliaria, petechial rash, pruritic rash, pustular rash, vesiculobullous rash, seborrhea, skin atrophy, skin hypertrophy,

skin striae, sweating decreased. Special senses - Frequent: abnormality of accommodation, mydriasis, taste perversion; Infrequent: conjunctivitis, diplopia, dry eyes, otitis media, parosmia, photophobia, taste loss; Rare: blepharitis, cataract, chromatopsia, conjunctival edema, corneal lesion, deafness, exophthalmos, eye hemorrhage, angle-closure glaucoma, retinal hemorrhage, subconjunctival hemorrhage, hyperacusis, keratitis, labyrinthitis, miosis, papilledema, decreased pupillary reflex, otitis externa, scleritis, uveitis, visual field defect.

Urogenital system - Frequent: albuminuria, urination impaired; Infrequent: amenorrhea,* cystitis, dysuria, hematuria, kidney calculus, kidney pain, leukorrhea,* menorrhagia,* metrorrhagia,* nocturia, breast pain, polyuria, pyuria, prostatic disorder (prostatitis, enlarged prostate, and prostate irritability),* urinary incontinence, urinary retention, urinary urgency, vaginal hemorrhage,* vaginitis*; Rare: abortion,* anuria, breast discharge, breast engorgement, balanitis,* breast enlargement, endometriosis,* female lactation,* fibrocystic breast, calcium crystalluria, cervicitis,* orchitis,* ovarian cyst,* bladder pain prolonged erection,* gynecomastia (male),* hypomenorrhea,* mastitis, menopause,* pyelonephritis, oliguria, salpingitis,* urolithiasis, uterine hemorrhage,* uterine spasm,* vaginal dryness.*

* Based on the number of men and women as appropriate.

6.2 Post-Marketing Experience Voluntary reports of other adverse reactions temporally associated with the use of venlafaxine have been received since market introduction. Because these reactions are reported from a population of uncertain size, it is not always possible to reliably estimate

their frequency or establish a causal relationship to drug exposure. These reports include the following reactions: agranulocytosis, anaphylaxis, aplastic anemia, catatonia, congenital anomalies, impaired coordination and balance, CPK increased, deep vein thrombophlebitis, delirium, Takotsubo cardiomyopathy, EKG abnormalities such as QT prolongation; cardiac arrhythmias including atrial fibrillation, supraventricular tachycardia, ventricular extrasystoles, and rare reports of ventricular fibrillation and ventricular tachycardia, including torsade de pointes; toxic epidermal necrolysis/Stevens-Johnson Syndrome, erythema multiforme extrapyramidal symptoms (including dyskinesia and tardive dyskinesia), angle-closure glaucoma, hemorrhad including extrapyramidal symptoms (including dyskinesia and tardive dyskinesia), angle-closure glaucoma, hemorrhad including eye and gastrointestinal bleeding), hepatic reactions (including GGT elevation; abnormalities of unspecified liver function tests; liver damage, necrosis, or failure; and fatty liver), interstitial lung disease, involuntary movements, LDH increased, neuroleptic alignant syndrome-like reactions (including a case of a 10-year-old who may have been taking methylphenidate, was treated and recovered), neutropenia, night sweats, pancreatitis, pancytopenia, panic, prolactin increased, renal failure, rhabdomyolysis, serotonin syndrome, shock-like electrical sensations or tinnitus (in some cases, subsequent to the discontinuation of venlafaxine or tapering of dose), and syndrome of inappropriate antidiuretic hormone secretion (usually in the elderly).

7 DRUG INTERACTIONS

A single dose of ethanol (0.5 g/kg) had no effect on the pharmacokinetics of venlafaxine or 0-desmethylvenlafaxine (0DV) when venlafaxine was administered at 150 mg/dav in 15 healthy male subjects. Additionally ventalaxine was administered at 150 mg/day in 15 healthy male subjects. Additionally, administration of ventalaxine in a stable regimen did not exaggerate the psychomotor and psychometric effects induced by ethanol in these same subjects when they were not receiving venlafaxine

7.2 Cimetidine Concomitant administration of cimetidine and venlafaxine in a steady-state study for both drugs resulted in inhibition of first-pass metabolism of venlafaxine in 18 healthy subjects. The oral clearance of venlafaxine was reduced by about 43%, and the exposure (AUC) and maximum concentration (C_{max}) of the drug were increased by about 60%. However, coadministration of cimetidine had no apparent effect on the pharmacokinetics of ODV, which is present in much greater quantity in the circulation than venlafaxine. The overall pharmacological activity of venlafaxine plus ODV is expected to increase only slightly, and no dosage adjustment should be necessary for most normal adults. However, for patients with pre-existing hypertension, and for elderly patients or patients with hepatic dysfunction, the interaction associated with the concomitant use of ventafaxine and cimetidine is not known and potentially could be more pronounced. Therefore, caution is advised with such patients.

Under steady-state conditions for venlafaxine administered at 150 mg/day, a single 10 mg dose of diazepam did not appear to affect the pharmacokinetics of either venlafaxine or ODV in 18 healthy male subjects. Venlafaxine also did not have any effect on the pharmacokinetics of diazepam or its active metabolite, desmethyldiazepam, or affect the psychomotor and psychometric effects induced by diazepam.

7.4 Haloperidol Venlafaxine administered under steady-state conditions at 150 mg/day in 24 healthy subjects decreased total oral-dose clearance (Cl/F) of a single 2 mg dose of haloperidol by 42%, which resulted in a 70% increase in haloperidol AUC. In addition, the haloperidol

increased 88% when coadministered with venlafaxine, but the haloperidol elimination half-life (t,a) was unchanged. The The steady-state pharmacokinetics of venlafaxine administered at 150 mg/day were not affected when a single 600 mg oral dose of

lithium was administered to 12 healthy male subjects, ODV also was unaffected. Venlafaxine had no effect on the pharmacokinetics of lithium (see also CNS-Active Drugs, below)

7.6 Drugs Highly Bound to Plasma ProteinsVenlafaxine is not highly bound to plasma proteins; therefore, administration of venlafaxine extended-release tablets to a patient taking another drug that is highly protein bound should not cause increased free concentrations of the other drug.

7.7 Drugs that Inhibit Cytochrome P450 Isoenzymes CYP2D6 Inhibitors: *In vitro* and *in vivo* studies indicate that venlafaxine is metabolized to its active metabolite, ODV, by CYP2D6 the isoenzyme that is responsible for the genetic polymorphism seen in the metabolism of many antidepressants. Therefore the potential exists for a drug interaction between drugs that inhibit CYP2D6-mediated metabolism of venlafaxine, reducing the metabolism of venlataxine to ODV, resulting in increased plasma concentrations of venlataxine and decreased concentrations of the active metabolist. CYP2D6 inhibitors such as quinidine would be expected to do this, but the effect would be similar to what is seen in patients who are genetically CYP2D6 poor metabolizers [see Clinical Pharmacology (12.3)]. Therefore, no dosage adjustment is

required when ventafaxine is coadministered with a CYP2D6 inhibitor. Ketoconazole: A pharmacokinetic study with ketoconazole 100 mg b.i.d. with a single dose of ventafaxine 50 mg in extensive metabolizers (EM; n=14) and 25 mg in poor metabolizers (PM;n=6) of CYP2D6 resulted in higher plasma concentrations of both venlafaxine and 0-desmethylvenlafaxine (ODV) in most subjects following administration of ketoconazole. Venlafaxine C_{max} increased by 26% in EM subjects and 48% in PM subjects. C_{max} values for ODV increased by 14% and 29% in EM and PM subjects,

respectively. enlafaxine AUC increased by 21% in EM subjects and 70% in PM subjects (range in PM's -2% to 206%), and AUC values for ODV increased by 23% and 33% in EM and PM (range in PM's -38% to 105%) subjects, respectively. Combined AUC's of venlafaxine and ODV increased on average by approximately 23% in EM's and 53% in PM's, (range in PM's 4% to 134%). Concomitant use of CYP3A4 inhibitors and venlafaxine may increase levels of venlafaxine and ODV. Therefore, caution is advised if a patient's therapy includes a CYP3A4 inhibitor and venlafaxine concomitantly.

7.8 Drugs Metabolized by Cytochrome P450 Isoenzymes

In vitro studies indicate that venlafaxine is a relatively weak inhibitor of CYP2D6. These findings have been confirmed in a clinical drug interaction study comparing the effect of venlafaxine with that of fluoxetine on the CYP2D6-mediated metabolism of

dextromethorphan to dextrorphan. Imipramine - Venlafaxine did not affect the pharmacokinetics of imipramine and 2-0H-imipramine. However, desipramine AUC $C_{\rm max}$, and $C_{\rm min}$ increased by about 35% in the presence of venlafaxine. The 2-OH-desipramine AUC's increased by at least 2.5 fold (with venlafaxine 37.5 mg q12h) and by 4.5 fold (with venlafaxine 75 mg q12h). Imipramine did not affect the pharmacokinetics of venlafaxine and ODV. The clinical significance of elevated 2-OH-desipramine levels is unknown.

Metoprolol - Concomitant administration of venlafaxine (50 mg every 8 hours for 5 days) and metoprolol (100 mg every 24 hours for 5 days) to 18 healthy male subjects in a pharmacokinetic interaction study for both drugs resulted in an increase of plasma concentrations of metoprolol by approximately 30-40% without aftering the plasma concentrations of its active metabolite, α -hydroxymetoprolol. Metoprolol did not alter the pharmacokinetic profile of venlafaxine or its active metabolite

0-desmethylvenlafaxine Venlafaxine appeared to reduce the blood pressure lowering effect of metoprolol in this study. The clinical relevance of this finding

or hypertensive patients is unknown. Caution should be exercised with co-administration of venlafaxine and metoprolol. Venlafaxine treatment has been associated with dose-related increases in blood pressure in some patients. It is recommended that patients receiving venlafaxine extended-release tablets have regular monitoring of blood pressure [see Warnings and Precautions

Risperidone - Venlafaxine administered under steady-state conditions at 150 mg/day slightly inhibited the CYP2D6-mediated metabolism of risperidone (administered as a single 1 mg oral dose) to its active metabolism (»-phydroxyrisperidone, resulting in an approximate 32% increase in risperidone AUC. However, venlafaxine coadministration did not significantly alter the pharmacokinetic profile of the total active moiety (risperidone plus 9-hydroxyrisperidone) CYP3A4

Venlafaxine did not inhibit CYP3A4 in vitro. This finding was confirmed in vivo by clinical drug interaction studies in which venlafaxine did not inhibit the metabolism of several CYP3A4 substrates, including alprazolam, diazepam, and terfenadine. Indinavir - In a study of 9 healthy volunteers, venlafaxine administered under steady-state conditions at 150 mg/day resulted in a 28% decrease in the AUC of a single 800 mg oral dose of indinavir and a 36% decrease in indinavir $C_{\text{max.}}$ Indinavir did not affect the pharmacokinetics of venlafaxine and ODV. The clinical significance of this finding is unknown

Venlafaxine did not inhibit CYP1A2 in vitro. This finding was confirmed in vivo by a clinical drug interaction study in which venlafaxine did not inhibit the metabolism of caffeine, a CYP1A2 substrate

CYP2C9 faxine did not inhibit CYP2C9 in vitro. In vivo, venlafaxine 75 mg by mouth every 12 hours did not alter the pharmacokinetics of a single 500 mg dose of tolbutamide or the CYP2C9 mediated formation of 4-hydroxy-tolbutamid

Venlafaxine did not inhibit the metabolism of diazepam, which is partially metabolized by CYP2C19 (see Diazepam above)

7.9 Monoamine Oxidase Inhibitors (MAOIs) [see Dosage and Administration (2.6 and 2.7), Contraindications (4.1), and Warnings and Precautions (5.2).]

7.10 Serotonergic Drugs [see Dosage and Administration (2.6 and 2.7), Contraindications (4.1), and Warnings and Precautions (5.2).]

sis (e.g., NSAID's, Aspirin, and Wa 7.11 Drugs that Interfere with He Serotonin release by platelets plays an important role in hemostasis. Epidemiological studies of the case-control and cohort

design that have demonstrated an association between use of psychotropic drugs that interfere with serotonin reuptake and the occurrence of upper gastrointestinal bleeding. These studies have also shown that concurrent use of an NSAID or aspirin may potentiate this risk of bleeding. Altered anticoagulant effects, including increased bleeding, have been reported when SSRI's and Skill's are coadministered with warfarin. Patients receiving warfarin therapy should be carefully monitored when venlata: extended-release tablets are initiated or discontinued. [see Warnings and Precautions (5.13).]

7.12 Electroconvulsive Therapy There are no clinical data establishing the benefit of electroconvulsive therapy combined with venlafaxine extended-release tablets

7.13 Postmarketing Spontaneous Drug Interaction Reports

here have been reports of elevated clozapine levels that were temporally associated with adverse reactions, including seizures, following the addition of venlafaxine. There have been reports of increases in prothrombin time, partial thromboplastin time, or INR when venlafaxine was given to

7.14 Drug-Laboratory Test Interactions False-positive urine immunoassay screening tests for phencyclidine (PCP) and amphetamine have been reported in patients

taking ventalaxine. This is due to lack of specificity of the screening tests. False positive test results may be expected for several days following discontinuation of ventafaxine therapy. Confirmatory tests, such as gas chromatography/mass spectrometry, will distinguish venlafaxine from PCP and amphetamine. 8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

patients receiving warfarin therapy.

Teratogenic Effects Pregnancy Category C

faxine did not cause malformations in offspring of rats or rabbits given doses up to 2.5 times (rat) or 4 times (rabbit) the maximum recommended human daily dose on a mg/m2 basis.

However, in rats, there was a decrease in pup weight, an increase in stillborn pups, and an increase in pup deaths during the first 5 days of lactation, when do sloing began during pregnancy and continued until wearing. The cause of these deaths is not known. These effects occurred at 2.5 times (mg/m²) the maximum human daily dose. The no effect dose for rat pup mortality was 0.25 times the human dose on a mg/m² basis. There are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly Non-Teratogenic Effects

Negrates exposed to veniafaxine hydrochloride extended-release capsules, other SNRIs (Serotonin and Noreninenhrine Reuntake Inhibitors), or SSRIs (Selective Serotonin Reuptake Inhibitors), late in the third trimester have developed complications requiring prolonged hospitalization, respiratory support, and tube feeding. Such complications can arise immediately upon delivery. Reported clinical findings have included respiratory distress, cyanosis, apnea, seizures, temperature instability, feeding difficulty, vomiting hypoglycemia, hypotonia, hypertonia, hyperteflexia, tremor, jitteriness, irritability, and constant crying. These features are consistent with either a direct toxic effect of SSRIs and SNRIs or, possibly, a drug discontinuation syndrome. It should be noted that, in some cases, the clinical picture is consistent with serotonin syndrome [see Warnings and Precautions (5.2)]. When treating a pregnant woman with venlafaxine extended-release tablets during the third trimester, the physician should carefully consider the potential risks and benefits of treatment [see Dosage and Administration (2)].

8.2 Labor and Delivery The effect of venlafaxine on labor and delivery in humans is unknown

8.3 Nursing Mothers

nlafaxine and ODV have been reported to be excreted in human milk. Because of the potential for serious adverse reactions in nursing infants from venlafaxine extended-release tablets, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother. 8.4 Pediatric Use

afety and effectiveness in the pediatric population have not been established [see BOXED WARNING and Warnings and Precaution (5.1). Two placebo-controlled trials in 766 pediatric patients with MDD and two placebo-controlled trials in another disorder in 793 pediatric patients have been conducted with venlafaxine hydrochloride extended-release capsules, and the data were not sufficient Anyone considering the use of venlafaxine extended-release tablets in a child or adolescent must balance the potential risks with

the clinical need. Although no studies have been designed to primarily assess impact of venlafaxine hydrochloride extended-release capsules on the growth, development, and maturation of children and adolescents, the studies that have been done suggest that venlataxine extended-release tablets may adversely affect weight and height [see Warnings and Precautions (5.7, 5.8, and 5.9]]. Should the decision be made to treat a pediatric patient with venlafaxine extended-release tablets, regular monitoring of weight and height is recommended during treatment, particularly if it is to be continued long term. The safety of venlataxine extended-release tablet treatment for pediatric patients has not been systematically assessed for chronic treatment longer than six months in duration. In the studies conducted in pediatric patients (ages 6-17), the occurrence of blood pressure and cholesterol increases considered

to be clinically relevant in pediatric patients was similar to that observed in adult patients. Consequently, the precautions for adults

Approximately 4% (14/357) and 2% (6/277) of patients treated with venlafaxine hydrochloride extended-release capsules in placebo-controlled premarketing major depressive disorder and Social Anxiety Disorder trials, respectively, were 65 years of age or over. Of 2,887 patients treated with venlafaxine hydrochloride immediate-release tablets in premarketing phase major depressive disorder studies, 12% (357) were 65 years of age or over. No overall differences in effectiveness or safety were observed between geriatric patients and younger patients, and other reported clinical experience generally has not identified differences in response between the elderly and younger patients. However, greater sensitivity of some older individuals cannot be ruled out. SSRIs and SNRIs, including venlafaxine hydrochloride extended-release capsules have been associated with cases of clinically significant hyponatremia in elderly patients, who may be at greater risk for this adverse reaction [see Warnings and Precautions (5.11)]. The pharmacokinetics of venlafaxine and ODV are not substantially altered in the elderly [see Clinical Pharmacology (12.3]]. No

nended for the elderly on the basis of age alone, although other clinical circu may be more common in the elderly, such as renal or hepatic impairment, may warrant a dose reduction [see Dosage and Administration (2.31). 8.6 Patients with Hepatic Impairment In patients with cirrhosis of the liver, the clearances of venlafaxine and its active metabolite (0DV) were decreased, thus prolonging the elimination half-lives of these substances. A large degree of intersubject variability was noted. [see Clinical Pharmacology

(12.3).1 A lower dose and individualization of dosing may be necessary [see Dosage and Administration (2.3)]. Venlafaxine extended

8.7 Patients with Renal Impairmen In patients with renal impairment (GFR = 10 to 70 mL/min), the clearances of venlafaxine and its active metabolites were decreased, thus prolonging the elimination half-lives of these substances [see Clinical Pharmacology (12.3)]. It is recommended that the total daily dose be reduced by 25% to 50% in patients with renal impairment. Because there was much individual variability in clearance between patients with renal impairment, individualization of dosage may be desirable in some patients. In patients undergoing hemodialysis, it is recommended that the total daily dose be reduced by 50%, [see Dosage and Administration (2,3).]

kine extended-release tablets, like all drugs effective in the treatment of major depressive disorder, should be used with caution in such patients. 9 DRUG ABUSE AND DEPENDENCE 9.1 Controlled Substance faxine extended-release tablets (venlafaxine hydrochloride) are not a controlled substance.

While venlafaxine has not been systematically studied in clinical trials for its potential for abuse, there was no indication of drugseeking behavior in the clinical trials. However, it is not possible to predict on the basis of premarketing experience the extent to which a CNS active drug will be misused, diverted, and/or abused once marketed. Consequently, physicians should carefully evaluate patients for history of drug abuse and follow such patients closely, observing them for signs of misuse or abuse of

In vitro studies revealed that venlafaxine has virtually no affinity for opiate, benzodiazepine, phencyclidine (PCP), or N-methyl-Daspartic acid (NMDA) receptors.

Venlafaxine was not found to have any significant CNS stimulant activity in rodents. In primate drug discrimination studies,

venlafaxine showed no significant stimulant or depressant abuse liability. ontinuation effects have been reported in patients receiving venlafaxine [see Dosage and Administration (2.4) and Warnings

and Precautions (5.5)].

Among the patients included in the premarketing evaluation of venlafaxine hydrochloride extended-release capsules, there were 2

10 OVERDOSAGE 10.1 Human Experience

reports of acute overdosage with venlafaxine hydrochloride extended-release capsules in major depressive disorder trials, either alone or in combination with other drugs. One patient took a combination of 6 g of venlafaxine hydrochloride extended-release capsules and 2.5 mg of lorazenam. This patient was hospitalized, treated symptomatically, and recovered without any untoward ects. The other patient took 2.85 g of venlafaxine hydrochloride extended-release capsules. This patient reported paresthesia of all four limbs but recovered without sequelae. There were no reports of acute overdose with venlafaxine hydrochloride extended-release capsules in Social Anxiety Disorder trials. Among the patients included in the premarketing evaluation with venlafaxine hydrochloride immediate-release tablets, there were

14 reports of acute overdose with venlafaxine, either alone or in combination with other drugs and/or alcohol. The majority of the reports involved ingestion in which the total dose of venlataxine taken was estimated to be no more than several-fold higher than the usual therapeutic dose. The 3 patients who took the highest doses were estimated to have ingested approximately 6.75 g, 2.75 g, and 2.5 g. The resultant peak plasma levels of venlafaxine for the latter 2 patients were 6.24 and 2.35 ug/mL, respectively, and the peak plasma levels of 0-desmethylvenlafaxine were 3.37 and 1.30 µg/mL, respectively. Plasma venlafaxine levels were not obtained for the patient who ingested 6.75 g of venlafaxine. All 14 patients recovered without sequelae. Most patients reported no symptoms. Among the remaining patients, somnolence was the most commonly reported symptom. The patient who ingested 2,75 gof ventafaxine was observed to have 2 generalized convulsions and a prolongation of QTc to 500 msec, compared with 405 msec at baseline. Mild sinus tachycardia was reported in 2 of the other patients.

In postmarketing experience, overdose with venlafaxine has occurred predominantly in combination with alcohol and/or other drugs. The most commonly reported reactions in overdosage include tachycardia, changes in level of consciousness (ranging from somnolence to coma), mydriasis, seizures, and vomiting. Electrocardiogram changes (e.g., prolongation of QT interval, bundle branch block, QRS prolongation), ventricular tachycardia, bradycardia, hypotension, rhabdomyolysis, vertigo, liver necrosis,

serotonin syndrome, and death have been reported. Published retrospective studies report that venlafaxine overdosage may be associated with an increased risk of fatal outcomes compared to that observed with SSRI antidepressant products, but lower than that for tricyclic antidepressants. Epidemi studies have shown that ventafaxine-treated patients have a higher pre-existing burden of suicide risk factors than SSRI-treated patients. The extent to which the finding of an increased risk of fatal outcomes can be attributed to the toxicity of ventafaxine in overdosage as opposed to some characteristic(s) of venlafaxine-treated patients is not clear. Prescriptions for venlafaxine lease tablets should be written for the smallest quantity of tablets consistent with good patient management, in order to reduce the risk of overdose.

10.2 Management of Overdosage Treatment should consist of those general measures employed in the management of overdosage with any antidepressant. Ensure an adequate airway oxygenation and ventilation Monitor cardiac rhythm and vital signs. General supportive and symptomatic measures are also recommended. Induction of emesis is not recommended. Gastric lavage with a large bore orogastric tube with appropriate airway protection, if needed, may be indicated if performed soon after ingestion or in symptomatic

Activated charcoal should be administered. Due to the large volume of distribution of this drug, forced diuresis, dialvsis, nemoperfusion, and exchange transfusion are unlikely to be of benefit. No specific antidotes for venlafaxine are know

In managing overdosage, consider the possibility of multiple drug involvement. The physician should consider contacting a poison control center for additional information on the treatment of any overdose. Telephone numbers for certified poison control centers are listed in the Physicians' Desk Reference® (PDR). 11 DESCRIPTION

afaxine extended-release tablets (venlafaxine hydrochloride) are extended-release tablets for oral administration that contain venlafaxine hydrochloride, a structurally novel antidepressant. Venlafaxine hydrochloride is a selective serotonin and norepinephrine reuptake inhibitor (SNRI). It is designated (R/S)-1-[2-(dimethylamino)-1-(4-methoxyphenyl)ethyl] cyclohexanol hydrochloride or (±)-1-[α-[(dimethylamino)methyl]-p-methoxybenzyl] cyclohexanol hydrochloride and has the empirical formula of $C_{17}H_{27}NO_2$ HCI. Its molecular weight is 313.87. The structural formula is shown below.

venlafaxine hydrochloride

axine hydrochloride is a off-white to white crystalline solid with a solubility of 572 mg/mL in water (adjusted to ionic strength of 0.2 M with sodium chloride), Its octanol:water (0.2 M sodium chloride) partition coefficient is 0.43.

Venlafaxine extended-release tablets are formulated as extended-release tablet for once-a-day oral administration. Venlafaxine extended-release tablets use osmotic pressure to deliver venlafaxine hydrochloride at a controlled rate over approximately 24 hours. The system, which resembles a conventional tablet in appearance, comprises an osmotically active core surrounded by a semipermeable membrane. The unitary tablet core is composed of the drug and excipients (including the osmotically active components). There is a precision-laser drilled orifice in the semipermeable membrane on the side of the tablet. In an aqueous environment, such as the gastrointestinal tract, water permeates through the membrane into the tablet core, causing the drug to dissolve and the osmotic components to expand. This expansion pushes the drug out through the orifice. The semipermeable membrane controls the rate at which water permeates into the tablet core, which in turn controls the rate of drug delivery. The controlled rate of drug delivery into the gastrointestinal lumen is thus independent of pH or gastrointestinal motility. The function of venlafaxine extended-release tablets depends on the existence of an osmotic gradient between the contents of the core and the fluid in the gastrointestinal tract. Since the osmotic gradient remains constant, drug delivery remains ess The biologically inert components of the tablet remain intact during gastrointestinal transit and are eliminated in the feces as an

Tablets contain venlafaxine hydrochloride, USP equivalent to 37.5 mg, 75 mg, 150 mg, or 225 mg venlafaxine. Inactive ingredients consist of mannitol, microcrystalline cellulose, povidone, polyethylene glycol, colloidal silicon dioxide, magnesium stearate, ellulose acetate, hypromellose, titanium dioxide and talc.

Each tablet strength also contains black iron oxide, hypromellose and propylene glycol as imprinting ink.

12 CLINICAL PHARMACOLOGY 12.1 Mechanism of Action

The mechanism of the antidepressant action of venlafaxine in humans is believed to be associated with its potentiation of neurotransmitter activity in the CNS. Preclinical studies have shown that venlafaxine and its active metabolite, O-desmethylvenlafaxine (ODV) are notent inhibitors of neuronal serotonin and noreninenhrine reuntake and weak inhibitors of

Venlafaxine and its active metabolite, O-desmethylvenlafaxine (ODV) have no significant affinity for muscarinic cholinergic, H₁histaminergic or α -adrenergic recentors in vitro. Pharmacologic activity at these recentors is hypothesized to be associated with he various anticholinergic, sedative, and cardiovascular effects seen with other psychotropic drugs. Venlafaxine and ODV do not possess monoamine oxidase (MAO) inhibitory activity.

Steady-state concentrations of venlafaxine and 0-desmethylyenlafaxine (ODV) in plasma are attained within 3 days of oral multiple dose therapy. Venlafaxine and ODV exhibited linear kinetics over the dose range of 75 to 450 mg/day. The mean ± SD apparent elimination half-life for venlafaxine and ODV after administration of 75 mg venlafaxine extended-release tablets under fed conditions were 10.7±3.2 hours and 12.5±3.0 hours respectively. Venlafaxine and ODV are minimally bound at therapeutic concentrations to plasma proteins (27% and 30%, respectively) Absorption and Distribution

VenIafaxine is well absorbed and extensively metabolized in the liver. ODV is the only major active metabolite. On the basis of mass balance studies, at least 92% of a single oral dose of venlafaxine is absorbed. The absolute bioavailability of venlafaxine is about 45%, Administration of 75 mg venlafaxine extended-release tablets under fed conditions resulted in mean ± SD venlafaxine adout 47%. Administration of 73 mg venidatalite extended-release tablets under fed conditions resulted in mean ± 3D Venidatalite (2... of 26.9 ± 13.4 ng/ml. and AUC of 153.6.3 ± 496.8 ng/hr/ml. T_{max} was 6.3 ± 2.3 hours. DDV mean ± SD C_{max} AUC, T_{max} flet administration of 75 mg venidataxine extended-release tablets under fed conditions were 97.9 ± 29.4 ng/mL, 2926.0 ± 746.1 ng-hr/ ml and 11.6 + 2.9 hours respectively

Administration of venlafaxine hydrochloride extended-release capsules (150 mg g24 hours), generally resulted in lower C (150 Administration of vientilatakine injoint controlled in the controlled region of the controlled in the T_{max}'s were 2 hours for venlafaxine and 3 hours for ODV). When equal daily doses of venlafaxine were administered as either an immediate release tablet or the extended-release form of venlafaxine, the exposure to both venlafaxine and ODV would be similar for the two treatments. Venlafaxine extended-release tablets would, therefore, provide a slower rate of absorption, but the same extent of absorption compared with the immediate release tablet.

Food did not affect the pharmacokinetic parameters ALIC C and T of venlafaxine or its active metabolite ODV after faxine extended-release tablets. Time of administration (AM vs PM) would not affect the pharmacokinetics of venlafaxine and ODV. Equal doses of venlafaxine extended-release tablets are bioequivalent to Effexor XR capsules when administered under fed conditions.

Metabolism and Excretion

Following absorption, venlafaxine undergoes extensive presystemic metabolism in the liver, primarily to ODV, but also to N-desmethylvenlafaxine, N,O-didesmethylvenlafaxine, and other minor metabolites. *In vitro* studies indicate that the formation of ODV is catalyzed by CYP2D6: this has been confirmed in a clinical study showing that patients with low CYP2D6 levels ("poor on our is catalyzed by CTF2DO, this has been commissed in a clinical study showing that patients with now circle evers (poor metabolizers) had increased levels of venlafaxine and reduced levels of ODV compared to people with normal CYP2DG ("extensive metabolizers"). The differences between the CYP2DG poor and extensive metabolizers, however, are not expected to be clinically important because the sum of venlafaxine and ODV is similar in the two groups and venlafaxine and ODV are pharmacologically approximately equiactive and equipotent.

Approximately 87% of a venlafaxine dose is recovered in the urine within 48 hours as unchanged venlafaxine (5%), unconjugated ODV (29%), conjugated ODV (26%), or other minor inactive metabolites (27%). Renal elimination of venlafaxine and its metabolites is thus the primary route of excretion. Special Populations

Age and Gender: A population pharmacokinetic analysis of 404 venlafaxine-treated patients from two studies involving both b.i.d. and t.i.d. regimens showed that dose-normalized trough plasma levels of either venlafaxine or ODV were unaltered by age or gender differences. Dosage adjustment based on the age or gender of a patient is generally not necessary [see Dosage and Extensive/Poor Metabolizers: Plasma concentrations of venlafaxine were higher in CYP2D6 poor metabolizers than extensive

metabolizers. Because the total exposure (AUC) of venlafaxine and ODV was similar in poor and extensive metabolizer groups, however, there is no need for different venlafaxine dosing regimens for these two groups. Liver Disease: In 9 subjects with hepatic cirrhosis, the pharmacokinetic disposition of both venlafaxine and ODV was significantly

altered after oral administration of venlafaxine. Venlafaxine elimination half-life was prolonged by about 30%, and clearance decreased by about 50% in cirrhotic subjects compared to normal subjects. ODV elimination half-life was prolonged by about 60%, and clearance decreased by about 30% in cirrhotic subjects compared to normal subjects. A large degree of intersubject bility was noted. Three patients with more severe cirrhosis had a more substantial decrease in venlafaxine clearance (about 90%) compared to normal subjects.

In a second study, venlafaxine was administered orally and intravenously in normal (n = 21) subjects, and in Child-Pugh A (n = 8) and Child-Pugh B (n = 11) subjects (mildly and moderately impaired, respectively). Venlafaxine oral bioavailability was increased 2 to 3 fold, oral elimination half-life was approximately twice as long and oral clearance was reduced by more than half, compared to normal subjects. In hepatically impaired subjects, ODV oral elimination half-life was prolonged by about 40%, while oral clearance for ODV was similar to that for normal subjects. A large degree of intersubject variability was noted.

Dosage adjustment is necessary in these hepatically impaired patients [see Dosage and Administration (2.3) and Use in Specific Renal Disease: In a renal impairment study, venlafaxine elimination half-life after oral administration was prolonged by about 50% and clearance was reduced by about 24% in renally impaired patients (GFR=10 to 70 mL/min), compared to normal subjects.

In dialysis patients, venlafaxine elimination half-life was prolonged by about 180% and clearance was reduced by about 57% compared to normal subjects. Similarly, ODV elimination half-life was prolonged by about 40% although clearance was unchanged in patients with renal impairment (GFR=10 to 70 mL/min) compared to normal subjects. In dialysis patients, ODV elimination half-life was prolonged by about 142% and clearance was reduced by about 56% compared to normal subjects. A large degree intersubject variability was noted. Dosage adjustment is necessary in these patients [see Dosage and Administration (2.3) and

13 NONCLINICAL TOXICOLOGY 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Venlafaxine was given by oral gavage to mice for 18 months at doses up to 120 mg/kg per day, which was 1.7 times the maximum recommended human dose on a mg/m² basis. Venlafaxine was also given to rats by oral gavage for 24 months at doses up to 120 mg/kg per day. In rats receiving the 120 mg/kg dose, plasma concentrations of venlafaxine at necropsy were 1 times (male rats) and 6 times (female rats) the plasma concentrations of patients receiving the maximum recommended human dose. Plasma levels of the 0-desmethyl metabolite were lower in rats than in patients receiving the maximum recommended dose. Tumors were not

Venlafaxine and the major human metabolite, 0-desmethylvenlafaxine (0DV), were not mutagenic in the Ames reverse mutation assay in Salmonella bacteria or the Chinese hamster ovary/HGPRT mammalian cell forward gene mutation assay. Venlafaxine was also not mutagenic or clastogenic in the in vitro BALB/c-3T3 mouse cell transformation assay, the sister chromatid exchange assay in cultured Chinese hamster ovary cells, or in the *in vivo* chromosomal aberration assay, in the marrow. ODV was not clastogenic in the *in vitro* Chinese hamster ovary cell chromosomal aberration assay, but elicited a clastogenic response in the *in* vivo chromosomal aberration assay in rat bone marrow. Impairment of Fertility

Reproduction and fertility studies in rats showed no effects on male or female fertility at oral doses of up to 2 times the maximum recommended human dose on a mg/m2 basis

14 CLINICAL STUDIES 14.1 Major Depressive Disorde

Use in Specific Populations (8.7).

The efficacy of venlafaxine hydrochloride extended-release capsules as a treatment for major depr in two placebo-controlled, short-term, flexible-dose studies in adult outpatients meeting DSM-III-R or DSM-IV criteria for major depressive disorder.

A 12-week study utilizing venlafaxine hydrochloride extended-release capsules doses in a range 75 to 150 mg/day (mean dose for completers was 136 mg/day) and an 8-week study utilizing venlafaxine hydrochloride extended-release capsules doses in a range 75 to 225 mg/day (mean dose for completers was 177 mg/day) both demonstrated superiority of venlafaxine hydrochloride ded-release capsules over placebo on the HAM-D total score, HAM-D Depressed Mood Item, the MADRS total score, the

Clinical Global Impressions (CGI) Severity of Illness item, and the CGI Global Improvement item. In both studies, venlafaxine hydrochloride extended-release capsules were also significantly better than placebo for certain factors of the HAM-D, including he anxiety/somatization factor, the cognitive disturbance factor, and the retardation factor, as well as for the psychic anxiety scor A 4-week study of inpatients meeting DSM-III-R criteria for major depressive disorder with melancholia utilizing venlafaxine hydrochloride immediate-release tablets in a range of 150 to 375 mg/day (t.i.d. schedule) demonstrated superiority of venlafaxine hydrochloride immediate-release tablets over placebo. The mean dose in completers was 350 mg/day.

Examination of gender subsets of the population studied did not reveal any differential responsiveness on the basis of gender. In one longer-term study, adult outpatients meeting DSM-IV criteria for major depressive disorder who had responded during an 8-week open trial on venlafaxine hydrochloride extended-release capsules (75, 150, or 225 mg, qAM) were randomized to continuation of their same venlafaxine hydrochloride extended-release capsules dose or to placebo, for up to 26 weeks of observation for relapse.

Response during the open phase was defined as a CGI Severity of Illness item score of <3 and a HAM-D-21 total score of <10. nesponse during the open prizes was defined as a dost severity of miness tent score or Saint a new-0-21 total score or STO at the day 56 evaluation. Relapse during the double-blind phase was defined as follows: (1) a regarance of major depressive disorder as defined by DSM-IV criteria and a CGI Severity of Illness item score of ≥4 (moderately ill), (2) 2 consecutive CGI Severity of Illness item scores of >4 or (3) a final CGI Severity of Illness item score of >4 for any patient who withdrew from the study for on miless term scores of 24, or (5) a final coal operancy or miless item score of 24 for any patient who windrew from the study for any reason. Patients receiving continued venlafaxine hydrochloride extended-release capsules treatment experienced significantly lower relapse rates over the subsequent 26 weeks compared with those receiving placebo.

In a second longer-term trial, adult outpatients meeting DSM-III-R criteria for major depressive disorder, recurrent type, who had responded (HAM-D-21 total score ≤12 at the day 56 evaluation) and continued to be improved [defined as the following criteria being met for days 56 through 180: (1) no HAM-D-21 total score ≥20; (2) no more than 2 HAM-D-21 total scores >10, and (3) no single CGI Severity of Illness item score ≥4 (moderately ill)] during an initial 26 weeks of treatment on venlafaxine hydrochloride immediate-release tablets (100 to 200 mg/day, on a b.i.d. schedule) were randomized to continuation of their same dose of venlafaxine hydrochloride immediate-release tablets or to placebo. The follow-up period to observe patients for relapse, defined as a CGI Severity of Illness item score \ge 4, was for up to 52 weeks. Patients receiving continued treatment with venlafaxine hydrochloride immediate-release tablets experienced significantly lower relapse rates over the subsequent 52 weeks compared

14.2 Social Anxiety Disorder (Social Phobia)

16 HOW SUPPLIED/STORAGE AND HANDLING

The efficacy of venlafaxine hydrochloride extended-release capsules as a treatment for Social Anxiety Disorder (also known as Social Phobia) was established in two double-blind, parallel group, 12-week, multicenter, placebo-controlled, flexible-dose studies in adult outpatients meeting DSM-IV criteria for Social Anxiety Disorder. Patients received doses in a range of 75 to 225 mg/day. Efficacy was assessed with the Liebowitz Social Anxiety Scale (LSAS). In these two trials, venlafaxine hydrochloride extendedrelease capsules were significantly more effective than placebo on change from baseline to endpoint on the LSAS total score. Examination of subsets of the population studied did not reveal any differential responsiveness on the basis of gender. There was

enlafaxine extended-release tablets 37.5 mg are white to off white, film coated, round biconvex tablets printed with "392" in black ink. They are supplied as follows: Unit of Use Bottles of 30 Tablets NDC 31722-123-30 Unit of Use Bottles of 90 Tablets NDC 31722-123-90

Venlafaxine extended-release tablets 75 mg are white to off white, film coated, round biconvex tablets printed with "393" in black ink. They are supplied as follows:
Unit of Use Bottles of 30 Tablets NDC 31722-124-30

insufficient information to determine the effect of age or race on outcome in these studies.

Unit of Use Bottles of 90 Tablets NDC 31722-124-90 Venifation extended-release tablets 150 mg are white to off white, film coated, round biconvex tablets printed with "394" in black ink. They are supplied as follows:

Unit of Use Bottles of 90 Tablets NDC 31722-125-90

Venlafaxine extended-release tablets 225 mg are white to off white, film coated, round biconvex tablets printed with "395" in black

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide). Prescribers or other health professionals should inform patients, their families, and their caregivers about the benefits and risks associated with treatment with venlafaxine extended-release tablets and should counsel them in its appropriate use. A patient Medication Guide about "Antidepressant Medicines, Depression and Other Serious Mental Illness, and Suicidal Thoughts or Actions" is available for venlafaxine extended-release tablets. The prescriber or health professional should instruct patients, their families, and their caregivers to read the Medication Guide and should assist them in understanding its contents. Patients should be given the opportunity to discuss the contents of the Medication Guide and to obtain answers to any questions they may have. The

complete text of the Medication Guide is reprinted at the end of this document.

Patients should be advised of the following issues and asked to alert their prescriber if these occur while taking venlafaxine 17.1 Clinical Worsening and Suicide Risk

professional, especially if they are severe, abrupt in onset, or were not part of the patient's presenting symptoms. Symptoms such as these may be associated with an increased risk for suicidal thinking and behavior and indicate a need for very close mon and possibly changes in the medication. 17.2 Interference with Cognitive and Motor Performance Clinical studies were performed to examine the effects of venlafaxine on behavioral performance of healthy individuals. The

results revealed no clinically significant impairment of psychomotor, cognitive, or complex behavior performance. However, since any psychoactive drug may impair judgment, thinking, or motor skills, patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that venlafaxine therapy does not adversely affect their ability

tablets and triptans, tricyclic antidepressants, fentanyl, lithium, tramadol, amphetamines, tryptophan, buspirone, and St. John's Wort supplements or other serotonergic agents [see Warnings and Precautions (5.2) and Drug Interactions (7.10)]. Patients should be cautioned about the concomitant use of venlafaxine extended-release tablets and NSAID's, aspirin, warfaring or other drugs that affect coagulation since combined use of psychotropic drugs that interfere with serotonin reuptake and these

Patients should be cautioned about the risk of serotonin syndrome with the concomitant use of venlafaxine extended-release

Although venlafaxine has not been shown to increase the impairment of mental and motor skills caused by alcohol, patients should be advised to avoid alcohol while taking venlafaxine 17.5 Allergic Reactions

17.7 Nursing Patients should be advised to notify their physician if they are breast-feeding an infant.

glaucoma, when diagnosed, can be treated definitively with iridectomy. Open-angle glaucoma is not a risk factor for angle closure glaucoma. Patients may wish to be examined to determine whether they are susceptible to angle closure, and have a prophylactic procedure (e.g., iridectomy), if they are susceptible. [see Warnings and Precautions (5.4)]

17.9 Sexual Dysfunctio Advise patient that use of venlafaxine extended-release tablets may cause symptoms of sexual dysfunction in both male and female patients. Inform patients that they should discuss any changes in sexual function and potential management strategies with

Manufactured by: Central Islin NY 11722 Manufactured for:

Medication Guide Venlafaxine Extended-Release Tablets

Antidepressant Medicines, Depression and Other Serious Mental Illnesses, and Suicidal Thoughts or Actions Read the Medication Guide that comes with your or your family member's antidepressant medicine. This Medication Guide is only about the risk of suicidal thoughts and actions with antidepressant medicines. Talk to your, or your family member's, healthcare

provider about: all risks and benefits of treatment with antidepressant medicines

all treatment choices for depression or other serious mental illness Who should not take venlafaxine extended-release tablets? take a monoamine oxidase inhibitor (MAOI). Ask your healthcare provider or pharmacist if you are not sure if you take an MAOI, including the antibiotic linezolid.

Do not take an MAOI within 7 days of stopping venlafaxine extended-release tablets unless directed to do so by your physician Do not start venlafaxine extended-release tablets if you stopped taking an MAOI in the last 2 weeks unless directed to do so by your physician.

within the first few months of treatment. 2. Depression and other serious mental illnesses are the most important causes of suicidal thoughts and actions. Some people may have a particularly high risk of having suicidal thoughts or actions. These include people who have (or have a

Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is very important when an antidepressant medicine is started or when the dose is changed. Call the healthcare provider right away to report new or sudden changes in mood, behavior, thoughts, or feelings.

Keep all follow-up visits with the healthcare provider as scheduled. Call the healthcare provider between visits as needed,

especially if you have concerns about symptoms. Call a healthcare provider right away if you or your family member has any of the following symptoms, especially if they are new, worse, or worry you:

 new or worse depression · other unusual changes in behavior or mood

 new or worse anxiety Visual Problems: Eye pain, change in vision, swelling or redness around eye · feeling very agitated or restless

· acting aggressive, being angry, or violent

What else do I need to know about antidepressant medicines?

Never stop an antidepressant medicine without first talking to a healthcare provider. Stopping an antidepressant medicine suddenly can cause other symptom Visual Problems. Only some people are at risk for these problems. You may want to undergo an eye examination to see if you are at risk and receive preventative treatment if you are.

treatment choices with the healthcare provider, not just the use of antidepressants. Antidepressant medicines have other side effects. Talk to the healthcare provider about the side effects of the medicine

prescribed for you or your family member. ressant medicines can interact with other medicines. Know all of the medicines that you or your family member takes. Keep a list of all medicines to show the healthcare provider. Do not start new medicines without first checking with vour healthcare provider. Not all antidepressant medicines prescribed for children are FDA approved for use in children. Talk to your child's

healthcare provider for more information. Sexual problems (dysfunction). Taking serotonin and norepinephrine reuptake inhibitors (SNRIs), including venlafaxine extended-release tablets, may cause sexual problems.

Delayed ejaculation or inability to have an ejaculation

Decreased sex drive

Problems getting or keeping an erection mptoms in females may include: Decreased sex drive

Delayed orgasm or inability to have an orgasm Talk to your healthcare provider if you develop any changes in your sexual function or if you have any questions or concerns about sexual problems during treatment with venlafaxine extended-release tablets. There may be treatments your healthcare provider can suggest.

Medication Guide available at http://camberpharma.com/medication-guides This Medication Guide has been approved by the U.S. Food and Drug Administration for all antidepressants

Manufactured for:

4296 Package Insert for Venlafaxine Extended-Release Tablets (Ascent-Camber) 126-10-2021.indd 2

10/19/21 12:38 PM

Unit of Use Bottles of 30 Tablets NDC 31722-125-30 ink They are supplied as follows: Unit of Use Bottles of 30 Tablets NDC 31722-126-30 Unit of Use Bottles of 90 Tablets NDC 31722-126-90

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. Protect from moisture and humidity

extended-release tablets.

Patients, their families, and their caregivers should be encouraged to be alert to the emergence of anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, mania, other unusual changes in behavior, worsening of depression, and suicidal ideation, especially early during antidepressant treatment and when the does is adjusted up or down. Families and caregivers of patients should be advised to look for the emergence of such symptoms on a day-to-day basis, since changes may be abrupt. Such symptoms should be reported to the patient's prescriber or health

17.3 Concomitant Medication
Patients should be advised to inform their physicians if they are taking, or plan to take, any prescription or over-the-counter drugs,

agents has been associated with an increased risk of bleeding [see Warnings and Precautions (5.13) and Drug Interactions (7.11)].

Patients should be advised to notify their physician if they develop a rash, hives, or a related allergic phenomenor Patients should be advised to notify their physician if they become pregnant or intend to become pregnant during therapy.

including herbal preparations and nutritional supplements, since there is a potential for interactions.

17.8 Angle Closure Glaucoma Patients should be advised that taking venlafaxine can cause mild pupillary dilation, which in susceptible individuals, can lead to an episode of angle closure glaucoma. Pre-existing glaucoma is almost always open-angle glaucoma because angle closure

their healthcare provider [see Warnings and Precautions (5.18)]

Camber Pharmaceuticals, Inc Piscataway, NJ 08854

What is the most important information I should know about antidepressant medicines, depression and other serious mental illnesses, and suicidal thoughts or actions? sant medicines may increase suicidal thoughts or actions in some children, teenagers, and young adults

amily history of) bipolar illness (also called manic-depressive illness) or suicidal thoughts or actions. 3. How can I watch for and try to prevent suicidal thoughts and actions in myself or a family men

· thoughts about suicide or dying · acting on dangerous impulses attempts to commit suicide · an extreme increase in activity and talking (mania)

· panic attacks · trouble sleeping (insomnia) new or worse irritability

Antidepressants are medicines used to treat depression and other illnesses. It is important to discuss all the risks of treating depression and also the risks of not treating it. Patients and their families or other caregivers should discuss all

Symptoms in males may include:

Call your doctor for medical advice about side effects, You may report side effects to FDA at 1-800-FDA-1088.

Manufactured by: Central Islip, NY 11722

Camber Pharmaceuticals, Inc Piscataway, NJ 08854